

Anti-VIP Receptor 1 Picoband Antibody
Catalog # ABO13028

Specification

Anti-VIP Receptor 1 Picoband Antibody - Product Information

Application	WB
Primary Accession	P32241
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for VIP Receptor 1 detection. Tested with WB in Human;Mouse;Rat.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-VIP Receptor 1 Picoband Antibody - Additional Information

Gene ID 7433

Other Names

Vasoactive intestinal polypeptide receptor 1, VIP-R-1, Pituitary adenylate cyclase-activating polypeptide type II receptor, PACAP type II receptor, PACAP-R-2, PACAP-R2, VPAC1, VIPR1

Application Details

Western blot, 0.1-0.5 µg/ml

Subcellular Localization

Cell membrane; Multi-pass membrane protein.

Tissue Specificity

In lung, HT-29 colonic epithelial cells, Raji B-lymphoblasts. Lesser extent in brain, heart, kidney, liver and placenta. Not expressed in CD4+ or CD8+ T-cells. Expressed in the T-cell lines HARRIS, HuT 78, Jurkat and SUP-T1, but not in the T- cell lines Peer, MOLT-4, HSB and YT.

Contents

Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg NaN₃.

Immunogen

A synthetic peptide corresponding to a sequence of human VIP Receptor 1 (QAELRRKWRRWHLQGVLGWNPKYRH).

Cross Reactivity

No cross reactivity with other proteins.

Storage

At -20°C; for one year. After r°Constitution, at 4°C; for one month. It°Can also be

aliquotted and stored frozen at -20°C ; for a longer time. Avoid repeated freezing and thawing.

Anti-VIP Receptor 1 Picoband Antibody - Protein Information

Name VIPR1

Function

This is a receptor for VIP. The activity of this receptor is mediated by G proteins which activate adenylyl cyclase. The affinity is $\text{VIP} = \text{PACAP-27} > \text{PACAP-38}$.

Cellular Location

Cell membrane; Multi-pass membrane protein.

Tissue Location

In lung, HT-29 colonic epithelial cells, Raji B- lymphoblasts. Lesser extent in brain, heart, kidney, liver and placenta. Not expressed in CD4+ or CD8+ T-cells. Expressed in the T- cell lines HARRIS, HuT 78, Jurkat and SUP-T1, but not in the T-cell lines Peer, MOLT-4, HSB and YT.

Anti-VIP Receptor 1 Picoband Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-VIP Receptor 1 Picoband Antibody - Images




Figure 1. Western blot analysis of VIP Receptor 1 using anti-VIP Receptor 1 antibody (ABO13028).

Anti-VIP Receptor 1 Picoband Antibody - Background

VIPR1 (Vasoactive intestinal polypeptide receptor 1), also known as VIPR, HVR1, is a protein that in humans is encoded by the VIPR1 gene. Distinct subsets of neural, respiratory, gastrointestinal, and immune cells bear specific high-affinity G protein-coupled receptors for VIP, such as VIPR1. The VIPR1 gene is mapped on 3p22.1. The VIPR1 gene was found to span approximately 22 kb and to be comprised of 13 exons (ranging from 42 to 1,400 bp) and 12 introns (ranging from 0.3 to 6.1 kb). One encodes a VIP receptor consisting of 460 amino acids and having 7 putative transmembrane domains, as do other G protein-coupled receptors. Patients with idiopathic achalasia show a significant difference in the distribution of SNPs affecting VIPR1.