

Anti-RECK Picoband Antibody
Catalog # ABO13052**Specification****Anti-RECK Picoband Antibody - Product Information**

Application	WB
Primary Accession	O95980
Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for RECK detection. Tested with WB in Human;Mouse.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-RECK Picoband Antibody - Additional Information

Gene ID 8434

Other Names

Reversion-inducing cysteine-rich protein with Kazal motifs, hRECK, Suppressor of tumorigenicity 15 protein, RECK, ST15

Application Details

Western blot, 0.1-0.5 µg/ml

Subcellular Localization

Cell membrane; Lipid-anchor, GPI-anchor.

Tissue Specificity

Expressed in various tissues and untransformed cells. It is undetectable in tumor-derived cell lines and oncogenically transformed cells.

Contents

Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg NaN₃.

Immunogen

A synthetic peptide corresponding to a sequence of human RECK (NAQSDQGAMNDMKLWEKGSIKMPFINIPVLDIKKCQPEMWKAIA).

Cross Reactivity

No cross reactivity with other proteins.

Storage

At -20°C; for one year. After r°Constitution, at 4°C; for one month. It°Can also be aliquotted and stored frozen at -20°C; for a

longer time. Avoid repeated freezing and thawing.

Anti-RECK Picoband Antibody - Protein Information

Name RECK {ECO:0000303|PubMed:9789069, ECO:0000312|HGNC:HGNC:11345}

Function

Functions together with ADGRA2 to enable brain endothelial cells to selectively respond to Wnt7 signals (WNT7A or WNT7B) (PubMed: 28289266, PubMed: 30026314). Plays a key role in Wnt7-specific responses: required for central nervous system (CNS) angiogenesis and blood-brain barrier regulation (By similarity). Acts as a Wnt7-specific coactivator of canonical Wnt signaling by decoding Wnt ligands: acts by interacting specifically with the disordered linker region of Wnt7, thereby conferring ligand selectivity for Wnt7 (PubMed: 30026314). ADGRA2 is then required to deliver RECK-bound Wnt7 to frizzled by assembling a higher-order RECK-ADGRA2-Fzd-LRP5-LRP6 complex (PubMed: 30026314). Also acts as a serine protease inhibitor: negatively regulates matrix metalloproteinase-9 (MMP9) by suppressing MMP9 secretion and by direct inhibition of its enzymatic activity (PubMed: 18194466, PubMed: 9789069). Also inhibits metalloproteinase activity of MMP2 and MMP14 (MT1-MMP) (PubMed: 9789069).

Cellular Location

Cell membrane; Lipid-anchor, GPI-anchor

Tissue Location


Expressed in various tissues and untransformed cells (PubMed:9789069). It is undetectable in tumor-derived cell lines and oncogenically transformed cells (PubMed:9789069)

Anti-RECK Picoband Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-RECK Picoband Antibody - Images






Figure 1. Western blot analysis of RECK using anti-RECK antibody (ABO13052).

Anti-RECK Picoband Antibody - Background

Reversion-inducing-cysteine-rich protein with kazal motifs, also known as RECK, is a human gene, thought to be a metastasis suppressor. The protein encoded by this gene is a cysteine-rich, extracellular protein with protease inhibitor-like domains whose expression is suppressed strongly in many tumors and cells transformed by various kinds of oncogenes. In normal cells, this membrane-anchored glycoprotein may serve as a negative regulator for matrix metalloproteinase-9, a key enzyme involved in tumor invasion and metastasis. Several transcript variants encoding different isoforms have been found for this gene.