

Anti-beta Tubulin Rabbit Monoclonal Antibody, HRP Conjugated

Catalog # ABO13101

Specification

Anti-beta Tubulin Rabbit Monoclonal Antibody, HRP Conjugated - Product Information

Application WB
Primary Accession Q13509
Host Rabbit Isotype Rabbit IgG

Reactivity Rat, Human, Mouse

Clonality Monoclonal Format Liquid

Description

Anti-beta Tubulin Rabbit Monoclonal Antibody, HRP Conjugated . Tested in WB application. This antibody reacts with Human, Mouse, Rat.

Anti-beta Tubulin Rabbit Monoclonal Antibody, HRP Conjugated - Additional Information

Gene ID 10381

Other Names

Tubulin beta-3 chain, Tubulin beta-4 chain, Tubulin beta-III, TUBB3, TUBB4

Calculated MW 50433 MW KDa

Application Details WB 1:5000-1:20000

Subcellular Localization Cytoplasm, cytoskeleton.

Tissue Specificity

Expression is primarily restricted to central and peripheral nervous system. Greatly increased expression in most cancerous tissues..

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human beta Tubulin

Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated



freeze-thaw cycles.

Anti-beta Tubulin Rabbit Monoclonal Antibody, HRP Conjugated - Protein Information

Name TUBB3

Synonyms TUBB4

Function

Tubulin is the major constituent of microtubules, a cylinder consisting of laterally associated linear protofilaments composed of alpha- and beta-tubulin heterodimers (PubMed:34996871). Microtubules grow by the addition of GTP-tubulin dimers to the microtubule end, where a stabilizing cap forms (PubMed:34996871). Below the cap, tubulin dimers are in GDP-bound state, owing to GTPase activity of alpha- tubulin (PubMed:34996871). TUBB3 plays a critical role in proper axon guidance and maintenance (PubMed:20074521). Binding of NTN1/Netrin-1 to its receptor UNC5C might cause dissociation of UNC5C from polymerized TUBB3 in microtubules and thereby lead to increased microtubule dynamics and axon repulsion (PubMed:28483977). Plays a role in dorsal root ganglion axon projection towards the spinal cord (PubMed:28483977).

Cellular Location

Cytoplasm, cytoskeleton. Cell projection, growth cone {ECO:0000250|UniProtKB:Q9ERD7}. Cell projection, lamellipodium {ECO:0000250|UniProtKB:Q9ERD7}. Cell projection, filopodium {ECO:0000250|UniProtKB:Q9ERD7}

Tissue Location

Expression is primarily restricted to central and peripheral nervous system. Greatly increased expression in most cancerous tissues.


Anti-beta Tubulin Rabbit Monoclonal Antibody, HRP Conjugated - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

Anti-beta Tubulin Rabbit Monoclonal Antibody, HRP Conjugated - Images





Western blot analysis of beta Tubulin expression in (1) Jurkat cell lysate, (2) Human kidney lysate, (3) 3T3 cell lysate, (4) Mouse brain lysate, (5) C6 cell lysate, (6) Rat heart lysate (H01857). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions.

After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-TUBB3 monoclonal antibody (Catalog # H01857) overnight at 4° C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit lgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for TUBB3