

## Anti-Phospho-Chk1 (S296) CHEK1 Rabbit Monoclonal Antibody Catalog # ABO13134

### Specification

---

#### Anti-Phospho-Chk1 (S296) CHEK1 Rabbit Monoclonal Antibody - Product Information

Application	WB
Primary Accession	<a href="#">O14757</a>
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid

#### Description

Anti-Phospho-Chk1 (S296) CHEK1 Rabbit Monoclonal Antibody . Tested in WB application. This antibody reacts with Human.

#### Anti-Phospho-Chk1 (S296) CHEK1 Rabbit Monoclonal Antibody - Additional Information

**Gene ID** 1111

#### Other Names

Serine/threonine-protein kinase Chk1, 2.7.11.1, CHK1 checkpoint homolog, Cell cycle checkpoint kinase, Checkpoint kinase-1, CHEK1, CHK1

#### Calculated MW

54434 MW KDa

#### Application Details

WB 1:500-1:2000

#### Subcellular Localization

Nucleus. Cytoplasm. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Nuclear export is mediated at least in part by XPO1/CRM1. Also localizes to the centrosome specifically during interphase, where it may protect centrosomal CDC2 kinase from inappropriate activation by cytoplasmic CDC25B.

#### Tissue Specificity

Expressed ubiquitously with the most abundant expression in thymus, testis, small intestine and colon..

#### Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

#### Immunogen

A synthesized peptide derived from human Phospho-Chk1 (S296)

#### Purification

Affinity-chromatography

Storage

**Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.****Anti-Phospho-Chk1 (S296) CHEK1 Rabbit Monoclonal Antibody - Protein Information**

Name CHEK1

Synonyms CHK1

**Function**

Serine/threonine-protein kinase which is required for checkpoint-mediated cell cycle arrest and activation of DNA repair in response to the presence of DNA damage or unreplicated DNA (PubMed: <a href="http://www.uniprot.org/citations/11535615" target="\_blank">11535615</a>, PubMed: <a href="http://www.uniprot.org/citations/12399544" target="\_blank">12399544</a>, PubMed: <a href="http://www.uniprot.org/citations/12446774" target="\_blank">12446774</a>, PubMed: <a href="http://www.uniprot.org/citations/14559997" target="\_blank">14559997</a>, PubMed: <a href="http://www.uniprot.org/citations/14988723" target="\_blank">14988723</a>, PubMed: <a href="http://www.uniprot.org/citations/15311285" target="\_blank">15311285</a>, PubMed: <a href="http://www.uniprot.org/citations/15650047" target="\_blank">15650047</a>, PubMed: <a href="http://www.uniprot.org/citations/15665856" target="\_blank">15665856</a>, PubMed: <a href="http://www.uniprot.org/citations/32357935" target="\_blank">32357935</a>). May also negatively regulate cell cycle progression during unperturbed cell cycles (PubMed: <a href="http://www.uniprot.org/citations/11535615" target="\_blank">11535615</a>, PubMed: <a href="http://www.uniprot.org/citations/12399544" target="\_blank">12399544</a>, PubMed: <a href="http://www.uniprot.org/citations/12446774" target="\_blank">12446774</a>, PubMed: <a href="http://www.uniprot.org/citations/14559997" target="\_blank">14559997</a>, PubMed: <a href="http://www.uniprot.org/citations/14988723" target="\_blank">14988723</a>, PubMed: <a href="http://www.uniprot.org/citations/15311285" target="\_blank">15311285</a>, PubMed: <a href="http://www.uniprot.org/citations/15650047" target="\_blank">15650047</a>, PubMed: <a href="http://www.uniprot.org/citations/15665856" target="\_blank">15665856</a>). This regulation is achieved by a number of mechanisms that together help to preserve the integrity of the genome (PubMed: <a href="http://www.uniprot.org/citations/11535615" target="\_blank">11535615</a>, PubMed: <a href="http://www.uniprot.org/citations/12399544" target="\_blank">12399544</a>, PubMed: <a href="http://www.uniprot.org/citations/12446774" target="\_blank">12446774</a>, PubMed: <a href="http://www.uniprot.org/citations/14559997" target="\_blank">14559997</a>, PubMed: <a href="http://www.uniprot.org/citations/14988723" target="\_blank">14988723</a>, PubMed: <a href="http://www.uniprot.org/citations/15311285" target="\_blank">15311285</a>, PubMed: <a href="http://www.uniprot.org/citations/15650047" target="\_blank">15650047</a>, PubMed: <a href="http://www.uniprot.org/citations/15665856" target="\_blank">15665856</a>). Recognizes the substrate consensus sequence [R-X-X- S/T] (PubMed: <a href="http://www.uniprot.org/citations/11535615" target="\_blank">11535615</a>, PubMed: <a href="http://www.uniprot.org/citations/12399544" target="\_blank">12399544</a>, PubMed: <a href="http://www.uniprot.org/citations/12446774" target="\_blank">12446774</a>, PubMed: <a href="http://www.uniprot.org/citations/14559997" target="\_blank">14559997</a>, PubMed: <a href="http://www.uniprot.org/citations/14988723" target="\_blank">14988723</a>, PubMed: <a href="http://www.uniprot.org/citations/15311285" target="\_blank">15311285</a>, PubMed: <a href="http://www.uniprot.org/citations/15650047" target="\_blank">15650047</a>, PubMed: <a href="http://www.uniprot.org/citations/15665856" target="\_blank">15665856</a>). Binds to and phosphorylates CDC25A, CDC25B and CDC25C (PubMed: <a href="http://www.uniprot.org/citations/12676583" target="\_blank">12676583</a>, PubMed: <a href="http://www.uniprot.org/citations/12676925" target="\_blank">12676925</a>, PubMed: <a href="http://www.uniprot.org/citations/12759351" target="\_blank">12759351</a>, PubMed: <a href="http://www.uniprot.org/citations/14559997" target="\_blank">14559997</a>, PubMed: <a href="http://www.uniprot.org/citations/14681206" target="\_blank">14681206</a>, PubMed: <a href="http://www.uniprot.org/citations/14681206" target="\_blank">14681206</a>, PubMed: <a href="http://www.uniprot.org/citations/14681206" target="\_blank">14681206</a>).

[19734889](http://www.uniprot.org/citations/19734889), PubMed: [9278511](http://www.uniprot.org/citations/9278511)). Phosphorylation of CDC25A at 'Ser-178' and 'Thr-507' and phosphorylation of CDC25C at 'Ser-216' creates binding sites for 14-3-3 proteins which inhibit CDC25A and CDC25C (PubMed: [9278511](http://www.uniprot.org/citations/9278511)). Phosphorylation of CDC25A at 'Ser-76', 'Ser-124', 'Ser-178', 'Ser-279' and 'Ser-293' promotes proteolysis of CDC25A (PubMed: [12676583](http://www.uniprot.org/citations/12676583), PubMed: [12676925](http://www.uniprot.org/citations/12676925), PubMed: [12759351](http://www.uniprot.org/citations/12759351), PubMed: [14681206](http://www.uniprot.org/citations/14681206), PubMed: [19734889](http://www.uniprot.org/citations/19734889), PubMed: [9278511](http://www.uniprot.org/citations/9278511)). Phosphorylation of CDC25A at 'Ser-76' primes the protein for subsequent phosphorylation at 'Ser-79', 'Ser-82' and 'Ser-88' by NEK11, which is required for polyubiquitination and degradation of CDC25A (PubMed: [19734889](http://www.uniprot.org/citations/19734889), PubMed: [20090422](http://www.uniprot.org/citations/20090422), PubMed: [9278511](http://www.uniprot.org/citations/9278511)). Inhibition of CDC25 leads to increased inhibitory tyrosine phosphorylation of CDK-cyclin complexes and blocks cell cycle progression (PubMed: [9278511](http://www.uniprot.org/citations/9278511)). Also phosphorylates NEK6 (PubMed: [18728393](http://www.uniprot.org/citations/18728393)). Binds to and phosphorylates RAD51 at 'Thr-309', which promotes the release of RAD51 from BRCA2 and enhances the association of RAD51 with chromatin, thereby promoting DNA repair by homologous recombination (PubMed: [15665856](http://www.uniprot.org/citations/15665856)). Phosphorylates multiple sites within the C-terminus of TP53, which promotes activation of TP53 by acetylation and promotes cell cycle arrest and suppression of cellular proliferation (PubMed: [10673501](http://www.uniprot.org/citations/10673501), PubMed: [15659650](http://www.uniprot.org/citations/15659650), PubMed: [16511572](http://www.uniprot.org/citations/16511572)). Also promotes repair of DNA cross-links through phosphorylation of FANCE (PubMed: [17296736](http://www.uniprot.org/citations/17296736)). Binds to and phosphorylates TLK1 at 'Ser-743', which prevents the TLK1-dependent phosphorylation of the chromatin assembly factor ASF1A (PubMed: [12660173](http://www.uniprot.org/citations/12660173), PubMed: [12955071](http://www.uniprot.org/citations/12955071)). This may enhance chromatin assembly both in the presence or absence of DNA damage (PubMed: [12660173](http://www.uniprot.org/citations/12660173), PubMed: [12955071](http://www.uniprot.org/citations/12955071)). May also play a role in replication fork maintenance through regulation of PCNA (PubMed: [18451105](http://www.uniprot.org/citations/18451105)). May regulate the transcription of genes that regulate cell-cycle progression through the phosphorylation of histones (By similarity). Phosphorylates histone H3.1 (to form H3T11ph), which leads to epigenetic inhibition of a subset of genes (By similarity). May also phosphorylate RB1 to promote its interaction with the E2F family of transcription factors and subsequent cell cycle arrest (PubMed: [17380128](http://www.uniprot.org/citations/17380128)). Phosphorylates SPRTN, promoting SPRTN recruitment to chromatin (PubMed: [31316063](http://www.uniprot.org/citations/31316063)). Reduces replication stress and activates the G2/M checkpoint, by phosphorylating and inactivating PABIR1/FAM122A and promoting the serine/threonine-protein phosphatase 2A-mediated dephosphorylation and stabilization of WEE1 levels and activity (PubMed: [33108758](http://www.uniprot.org/citations/33108758)).

### Cellular Location

Nucleus. Chromosome. Cytoplasm. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Note=Nuclear export is mediated at least in part by XPO1/CRM1 (PubMed:12676962). Also localizes to the centrosome specifically during interphase, where it may protect centrosomal

CDC2 kinase from inappropriate activation by cytoplasmic CDC25B (PubMed:15311285).  
Proteolytic cleavage at the C-terminus by SPRTN promotes removal from chromatin  
(PubMed:31316063)

#### Tissue Location


Expressed ubiquitously with the most abundant expression in thymus, testis, small intestine and colon

#### Anti-Phospho-Chk1 (S296) CHEK1 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### Anti-Phospho-Chk1 (S296) CHEK1 Rabbit Monoclonal Antibody - Images



Western blot analysis of Phospho-Chk1 (S296) expression in HEK293 cell lysate Treated with Calyculin.