

**Anti-Phospho-c-Jun (T91) Rabbit Monoclonal Antibody**  
Catalog # ABO13142**Specification****Anti-Phospho-c-Jun (T91) Rabbit Monoclonal Antibody - Product Information**

Application	WB, IP
Primary Accession	<a href="#">P05412</a>
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

**Description**

Anti-Phospho-c-Jun (T91) Rabbit Monoclonal Antibody . Tested in WB, IP applications. This antibody reacts with Human, Mouse, Rat.

**Anti-Phospho-c-Jun (T91) Rabbit Monoclonal Antibody - Additional Information**

**Gene ID** 3725

**Other Names**

Transcription factor Jun, Activator protein 1, AP1, Proto-oncogene c-Jun, Transcription factor AP-1 subunit Jun, V-jun avian sarcoma virus 17 oncogene homolog, p39, JUN

**Calculated MW**

35676 MW KDa

**Application Details**

WB 1:1000-1:2000<br>IP 1:20

**Subcellular Localization**

Nucleus.

**Contents**

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

**Immunogen**

A synthesized peptide derived from human c-Jun

**Purification**

Affinity-chromatography

**Storage**

**Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.**

**Anti-Phospho-c-Jun (T91) Rabbit Monoclonal Antibody - Protein Information**

**Name JUN****Function**

Transcription factor that recognizes and binds to the AP-1 consensus motif 5'-TGA[GC]TCA-3' (PubMed:<a href="http://www.uniprot.org/citations/10995748" target="\_blank">10995748</a>, PubMed:<a href="http://www.uniprot.org/citations/22083952" target="\_blank">22083952</a>). Heterodimerizes with proteins of the FOS family to form an AP-1 transcription complex, thereby enhancing its DNA binding activity to the AP-1 consensus sequence 5'-TGA[GC]TCA-3' and enhancing its transcriptional activity (By similarity). Together with FOSB, plays a role in activation-induced cell death of T cells by binding to the AP-1 promoter site of FASLG/CD95L, and inducing its transcription in response to activation of the TCR/CD3 signaling pathway (PubMed:<a href="http://www.uniprot.org/citations/12618758" target="\_blank">12618758</a>). Promotes activity of NR5A1 when phosphorylated by HIPK3 leading to increased steroidogenic gene expression upon cAMP signaling pathway stimulation (PubMed:<a href="http://www.uniprot.org/citations/17210646" target="\_blank">17210646</a>). Involved in activated KRAS-mediated transcriptional activation of USP28 in colorectal cancer (CRC) cells (PubMed:<a href="http://www.uniprot.org/citations/24623306" target="\_blank">24623306</a>). Binds to the USP28 promoter in colorectal cancer (CRC) cells (PubMed:<a href="http://www.uniprot.org/citations/24623306" target="\_blank">24623306</a>).

**Cellular Location**

Nucleus.

**Tissue Location**


Expressed in the developing and adult prostate and prostate cancer cells.

**Anti-Phospho-c-Jun (T91) Rabbit Monoclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Anti-Phospho-c-Jun (T91) Rabbit Monoclonal Antibody - Images**



Western blot analysis of Phospho-c-Jun (T91) expression in human brain lysate.