

Anti-Phospho-Cyclin E1 (T77) CCNE1 Rabbit Monoclonal Antibody
Catalog # ABO13184

Specification

Anti-Phospho-Cyclin E1 (T77) CCNE1 Rabbit Monoclonal Antibody - Product Information

Application	WB, IHC, IF, ICC
Primary Accession	P24864
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid

Description

Anti-Phospho-Cyclin E1 (T77) CCNE1 Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF applications. This antibody reacts with Human.

Anti-Phospho-Cyclin E1 (T77) CCNE1 Rabbit Monoclonal Antibody - Additional Information

Gene ID 898

Other Names

G1/S-specific cyclin-E1, CCNE1, CCNE

Calculated MW

47077 MW KDa

Application Details

WB 1:1000-1:2000
IHC 1:50-1:200
ICC/IF 1:50-1:200

Subcellular Localization

Nucleus.

Tissue Specificity

Highly expressed in testis and placenta. Low levels in bronchial epithelial cells..

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human Cyclin E1

Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated

freeze-thaw cycles.

Anti-Phospho-Cyclin E1 (T77) CCNE1 Rabbit Monoclonal Antibody - Protein Information

Name CCNE1

Synonyms CCNE

Function

Essential for the control of the cell cycle at the G1/S (start) transition.

Cellular Location

Nucleus.

Tissue Location


Highly expressed in testis and placenta. Low levels in bronchial epithelial cells.

Anti-Phospho-Cyclin E1 (T77) CCNE1 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Phospho-Cyclin E1 (T77) CCNE1 Rabbit Monoclonal Antibody - Images



Western blot analysis of Phospho-Cyclin E1 (T77) expression in (1) JAR cell treated with Lambda Phosphatase lysate; (2) JAR cell lysate.