

**Anti-AIF AIFM1 Rabbit Monoclonal Antibody**  
Catalog # ABO13226**Specification****Anti-AIF AIFM1 Rabbit Monoclonal Antibody - Product Information**

Application	WB, IHC, IF, ICC, IP, FC
Primary Accession	<a href="#">O95831</a>
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

**Description**

Anti-AIF AIFM1 Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

**Anti-AIF AIFM1 Rabbit Monoclonal Antibody - Additional Information**

**Gene ID** 9131

**Other Names**

Apoptosis-inducing factor 1, mitochondrial, 1.6.99.-, Programmed cell death protein 8, AIFM1 ([http://www.genenames.org/cgi-bin/gene\\_symbol\\_report?hgnc\\_id=8768](http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=8768) target="\_blank">HGNC:8768</a>), AIF, PDCD8

**Calculated MW**

66901 MW KDa

**Application Details**

WB 1:500-1:2000<br>IHC 1:100-1:500<br>ICC/IF 1:100-1:500<br>IP 1:50<br>FC 1:50

**Subcellular Localization**

Mitochondrion intermembrane space. Mitochondrion inner membrane. Cytoplasm. Nucleus. Cytoplasm, perinuclear region. Proteolytic cleavage during or just after translocation into the mitochondrial intermembrane space (IMS) results in the formation of an inner-membrane-anchored mature form (AIFmit). During apoptosis, further proteolytic processing leads to a mature form, which is confined to the mitochondrial IMS in a soluble form (AIFsol). AIFsol is released to the cytoplasm in response to specific death signals, and translocated to the nucleus, where it induces nuclear apoptosis. Colocalizes with EIF3G in the nucleus and perinuclear region.

**Tissue Specificity**

Detected in muscle and skin fibroblasts (at protein level). Isoform 5 is frequently down-regulated in human cancers..

**Contents**

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

**Immunogen**

A synthesized peptide derived from human AIF

#### Purification

Affinity-chromatography

Storage

**Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.**

### Anti-AIF AIFM1 Rabbit Monoclonal Antibody - Protein Information

Name AIFM1 ([HGNC:8768](#))

Synonyms AIF, PDCD8

#### Function

Functions both as NADH oxidoreductase and as regulator of apoptosis (PubMed:<a href="http://www.uniprot.org/citations/17094969" target="\_blank">17094969</a>, PubMed:<a href="http://www.uniprot.org/citations/20362274" target="\_blank">20362274</a>, PubMed:<a href="http://www.uniprot.org/citations/23217327" target="\_blank">23217327</a>, PubMed:<a href="http://www.uniprot.org/citations/33168626" target="\_blank">33168626</a>). In response to apoptotic stimuli, it is released from the mitochondrion intermembrane space into the cytosol and to the nucleus, where it functions as a proapoptotic factor in a caspase- independent pathway (PubMed:<a href="http://www.uniprot.org/citations/20362274" target="\_blank">20362274</a>). Release into the cytoplasm is mediated upon binding to poly-ADP-ribose chains (By similarity). The soluble form (AIFsol) found in the nucleus induces 'parthanatos' i.e. caspase-independent fragmentation of chromosomal DNA (PubMed:<a href="http://www.uniprot.org/citations/20362274" target="\_blank">20362274</a>). Binds to DNA in a sequence-independent manner (PubMed:<a href="http://www.uniprot.org/citations/27178839" target="\_blank">27178839</a>). Interacts with EIF3G, and thereby inhibits the EIF3 machinery and protein synthesis, and activates caspase-7 to amplify apoptosis (PubMed:<a href="http://www.uniprot.org/citations/17094969" target="\_blank">17094969</a>). Plays a critical role in caspase-independent, pyknotic cell death in hydrogen peroxide-exposed cells (PubMed:<a href="http://www.uniprot.org/citations/19418225" target="\_blank">19418225</a>). In contrast, participates in normal mitochondrial metabolism. Plays an important role in the regulation of respiratory chain biogenesis by interacting with CHCHD4 and controlling CHCHD4 mitochondrial import (PubMed:<a href="http://www.uniprot.org/citations/26004228" target="\_blank">26004228</a>).

#### Cellular Location

Mitochondrion intermembrane space. Mitochondrion inner membrane. Cytoplasm. Nucleus. Cytoplasm, perinuclear region. Note=Proteolytic cleavage during or just after translocation into the mitochondrial intermembrane space (IMS) results in the formation of an inner-membrane-anchored mature form (AIFmit). During apoptosis, further proteolytic processing leads to a mature form, which is confined to the mitochondrial IMS in a soluble form (AIFsol). AIFsol is released to the cytoplasm in response to specific death signals, and translocated to the nucleus, where it induces nuclear apoptosis (PubMed:15775970). Release into the cytoplasm is mediated upon binding to poly-ADP-ribose chains (By similarity) Translocation into the nucleus is promoted by interaction with (auto- poly-ADP-ribosylated) processed form of PARP1 (PubMed:33168626) Colocalizes with EIF3G in the nucleus and perinuclear region (PubMed:17094969). {ECO:0000250|UniProtKB:Q9Z0X1, ECO:0000269|PubMed:15775970, ECO:0000269|PubMed:17094969, ECO:0000269|PubMed:33168626} [Isoform 4]: Mitochondrion. Cytoplasm, cytosol. Note=In pro-apoptotic conditions, is released from mitochondria to cytosol in a calpain/cathepsin-dependent manner.

### Tissue Location


Expressed in all tested tissues (PubMed:16644725). Detected in muscle and skin fibroblasts (at protein level) (PubMed:23217327). Expressed in osteoblasts (at protein level) (PubMed:28842795). [Isoform 4]: Expressed in all tested tissues except brain.

### Anti-AIF AIFM1 Rabbit Monoclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


### Anti-AIF AIFM1 Rabbit Monoclonal Antibody - Images



Immunofluorescent analysis using the Antibody at 1:50 dilution.



Western blot analysis of AIF expression in K562 cell lysate.



Immunofluorescent analysis of HeLa cells, using AIF Antibody.