

**Anti-p63 TP63 Rabbit Monoclonal Antibody**  
Catalog # ABO13256**Specification****Anti-p63 TP63 Rabbit Monoclonal Antibody - Product Information**

|                   |                        |
|-------------------|------------------------|
| Application       | WB, IHC, IF, ICC, FC   |
| Primary Accession | <a href="#">O9H3D4</a> |
| Host              | Rabbit                 |
| Isotype           | Rabbit IgG             |
| Reactivity        | Rat, Human, Mouse      |
| Clonality         | Monoclonal             |
| Format            | Liquid                 |

**Description**

Anti-p63 TP63 Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

**Anti-p63 TP63 Rabbit Monoclonal Antibody - Additional Information**

**Gene ID** 8626

**Other Names**

Tumor protein 63, p63, Chronic ulcerative stomatitis protein, CUSP, Keratinocyte transcription factor KET, Transformation-related protein 63, TP63, Tumor protein p73-like, p73L, p40, p51, TP63, KET, P63, P73H, P73L, TP73L

**Calculated MW**

76785 MW KDa

**Application Details**

WB 1:500-1:2000<br>IHC 1:100-1:500<br>ICC/IF 1:50-1:200<br>FC 1:500

**Subcellular Localization**

Nucleus.

**Tissue Specificity**

Widely expressed, notably in heart, kidney, placenta, prostate, skeletal muscle, testis and thymus, although the precise isoform varies according to tissue type. Progenitor cell layers of skin, breast, eye and prostate express high levels of DeltaN-type isoforms. Isoform 10 is predominantly expressed in skin squamous cell carcinomas, but not in normal skin tissues..

**Contents**

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

**Immunogen**

A synthesized peptide derived from human p63

**Purification**

Affinity-chromatography

**Storage**

**Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.**

**Anti-p63 TP63 Rabbit Monoclonal Antibody - Protein Information**

**Name** TP63

**Synonyms** KET, P63, P73H, P73L, TP73L

**Function**

Acts as a sequence specific DNA binding transcriptional activator or repressor. The isoforms contain a varying set of transactivation and auto-regulating transactivation inhibiting domains thus showing an isoform specific activity. Isoform 2 activates RIPK4 transcription. May be required in conjunction with TP73/p73 for initiation of p53/TP53 dependent apoptosis in response to genotoxic insults and the presence of activated oncogenes. Involved in Notch signaling by probably inducing JAG1 and JAG2. Plays a role in the regulation of epithelial morphogenesis. The ratio of DeltaN-type and TA\*-type isoforms may govern the maintenance of epithelial stem cell compartments and regulate the initiation of epithelial stratification from the undifferentiated embryonal ectoderm. Required for limb formation from the apical ectodermal ridge. Activates transcription of the p21 promoter.

**Cellular Location**

Nucleus


**Tissue Location**


Widely expressed, notably in heart, kidney, placenta, prostate, skeletal muscle, testis and thymus, although the precise isoform varies according to tissue type. Progenitor cell layers of skin, breast, eye and prostate express high levels of DeltaN-type isoforms. Isoform 10 is predominantly expressed in skin squamous cell carcinomas, but not in normal skin tissues

**Anti-p63 TP63 Rabbit Monoclonal Antibody - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


**Anti-p63 TP63 Rabbit Monoclonal Antibody - Images**




All lanes use the Antibody at 1:2K dilution for 1 hour at room temperature.




All lanes use the Antibody at 1:2K dilution for 1 hour at room temperature.




All lanes use the Antibody at 1:2K dilution for 1 hour at room temperature.




All lanes use the Antibody at 1:2K dilution for 1 hour at room temperature.




All lanes use the Antibody at 1:2K dilution for 1 hour at room temperature.




All lanes use the Antibody at 1:2K dilution for 1 hour at room temperature.




Western blot analysis of p63 expression in A431 cell lysate.




Immunohistochemical analysis of paraffin-embedded Human thyroid cancer, using the Antibody at 1:500 dilution.




Immunohistochemical analysis of paraffin-embedded Human prostate cancer, using the Antibody at 1:500 dilution.



Immunohistochemical analysis of paraffin-embedded Human small cell lung cancer, using the Antibody at 1:500 dilution.



Immunohistochemical analysis of paraffin-embedded human breast, using p63 Antibody.



Immunofluorescent analysis of A431 cells, using p63 Antibody.