

## Anti-CD45 PTPRC Rabbit Monoclonal Antibody Catalog # ABO13351

### Specification

---

#### Anti-CD45 PTPRC Rabbit Monoclonal Antibody - Product Information

Application	WB, IHC, IF, ICC, FC
Primary Accession	<a href="#">P08575</a>
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid

#### Description

Anti-CD45 PTPRC Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, Flow Cytometry applications. This antibody reacts with Human.

#### Anti-CD45 PTPRC Rabbit Monoclonal Antibody - Additional Information

**Gene ID** 5788

#### Other Names

Receptor-type tyrosine-protein phosphatase C, 3.1.3.48, Leukocyte common antigen, L-CA, T200, CD45, PTPRC ([http://www.genenames.org/cgi-bin/gene\\_symbol\\_report?hgnc\\_id=9666](http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=9666) target="\_blank">HGNC:9666</a>), CD45

#### Calculated MW

147254 MW KDa

#### Application Details

WB 1:500-1:2000<br>IHC 1:50-1:200<br>ICC/IF 1:50-1:100<br>FC 1:20

#### Subcellular Localization

Membrane ; Single-pass type I membrane protein. Membrane raft. Colocalized with DPP4 in membrane rafts.

#### Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

#### Immunogen

A synthesized peptide derived from human CD45

#### Purification

Affinity-chromatography

#### Storage

**Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.**

## Anti-CD45 PTPRC Rabbit Monoclonal Antibody - Protein Information

Name PTPRC ([HGNC:9666](#))

Synonyms CD45

### Function

Protein tyrosine-protein phosphatase required for T-cell activation through the antigen receptor (PubMed:<a href="http://www.uniprot.org/citations/35767951" target="\_blank">35767951</a>). Acts as a positive regulator of T-cell coactivation upon binding to DPP4. The first PTPase domain has enzymatic activity, while the second one seems to affect the substrate specificity of the first one. Upon T-cell activation, recruits and dephosphorylates SKAP1 and FYN. Dephosphorylates LYN, and thereby modulates LYN activity (By similarity).

### Cellular Location

Cell membrane; Single-pass type I membrane protein. Membrane raft. Synapse. Note=Colocalized with DPP4 in membrane rafts.

### Tissue Location


Isoform 1: Detected in thymocytes. Isoform 2: Detected in thymocytes. Isoform 3: Detected in thymocytes. Isoform 4: Not detected in thymocytes. Isoform 5: Detected in thymocytes. Isoform 6: Not detected in thymocytes. Isoform 7: Detected in thymocytes Isoform 8: Not detected in thymocytes.


## Anti-CD45 PTPRC Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## Anti-CD45 PTPRC Rabbit Monoclonal Antibody - Images







Immunohistochemical analysis of paraffin-embedded Rat stomach, using the Antibody at 1:100 dilution.




Immunohistochemical analysis of paraffin-embedded Human placenta, using the Antibody at 1:100 dilution.




Immunohistochemical analysis of paraffin-embedded Mouse lung, using the Antibody at 1:100 dilution.



Immunohistochemical analysis of paraffin-embedded Mouse skin, using the Antibody at 1:100 dilution.



Western blot analysis of CD45 expression in Jurkat cell lysate.



Immunohistochemical analysis of paraffin-embedded human spleen, using CD45 Antibody.