

## Anti-CCR7 Rabbit Monoclonal Antibody Catalog # ABO13357

### Specification

---

#### Anti-CCR7 Rabbit Monoclonal Antibody - Product Information

Application	WB, IHC, IF, ICC, IP
Primary Accession	<a href="#">P32248</a>
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

#### Description

Anti-CCR7 Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, IP applications. This antibody reacts with Human, Mouse, Rat.

#### Anti-CCR7 Rabbit Monoclonal Antibody - Additional Information

**Gene ID** 1236

#### Other Names

C-C chemokine receptor type 7, C-C CKR-7, CC-CKR-7, CCR-7, BLR2, CDw197, Epstein-Barr virus-induced G-protein coupled receptor 1, EBI1, EBV-induced G-protein coupled receptor 1, MIP-3 beta receptor, CD197, CCR7, CMKBR7, EBI1, EVI1

#### Calculated MW

42874 MW KDa

#### Application Details

WB 1:500-1:2000<br>IHC 1:50-1:200<br>ICC/IF 1:50-1:200<br>IP 1:50

#### Subcellular Localization

Cell membrane; Multi-pass membrane protein.

#### Tissue Specificity

Expressed in various lymphoid tissues and activated B- and T-lymphocytes, strongly up-regulated in B-cells infected with Epstein-Barr virus and T-cells infected with herpesvirus 6 or 7.

#### Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150 mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5 mg/ml BSA.

#### Immunogen

A synthesized peptide derived from human CCR7

#### Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term

storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

## Anti-CCR7 Rabbit Monoclonal Antibody - Protein Information

**Name** CCR7

**Synonyms** CMKBR7, EBI1, EVI1

### Function

Receptor for the MIP-3-beta chemokine. Probable mediator of EBV effects on B-lymphocytes or of normal lymphocyte functions.

### Cellular Location

Cell membrane; Multi-pass membrane protein.

### Tissue Location


Expressed in various lymphoid tissues and activated B- and T-lymphocytes, strongly up-regulated in B-cells infected with Epstein-Barr virus and T-cells infected with herpesvirus 6 or 7

## Anti-CCR7 Rabbit Monoclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## Anti-CCR7 Rabbit Monoclonal Antibody - Images



Immunohistochemical analysis of paraffin-embedded human spleen, using CCR7 Antibody.



Immunofluorescent analysis of HeLa cells, using CCR7 Antibody .




Figure 1. Western blot analysis of CCR7 using anti-CCR7 antibody (M00390).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

- Lane 1: human MCF-7 whole cell lysates,
- Lane 2: human K562 whole cell lysates,
- Lane 3: rat spleen tissue lysates,
- Lane 4: rat lung tissue lysates,
- Lane 5: rat C6 whole cell lysates,
- Lane 6: mouse spleen tissue lysates,
- Lane 7: mouse lung tissue lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-CCR7 antigen affinity purified monoclonal antibody (Catalog # M00390) at 1:500 overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:500 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for CCR7 at approximately 43 kDa. The expected band size for CCR7 is at 43 kDa.