

**Anti-PGK1 Rabbit Monoclonal Antibody**  
Catalog # ABO13384**Specification****Anti-PGK1 Rabbit Monoclonal Antibody - Product Information**

Application	WB, IF, ICC, FC
Primary Accession	<a href="#">P00558</a>
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

**Description**

Anti-PGK1 Rabbit Monoclonal Antibody . Tested in WB, ICC/IF, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

**Anti-PGK1 Rabbit Monoclonal Antibody - Additional Information**

**Gene ID** 5230

**Other Names**

Phosphoglycerate kinase 1, 2.7.2.3, Cell migration-inducing gene 10 protein, Primer recognition protein 2, PRP 2, PGK1, PGKA

**Calculated MW**

44615 MW KDa

**Application Details**

WB 1:500-1:2000<br>ICC/IF 1:50-1:200<br>FC 1:50

**Subcellular Localization**

Cytoplasm.

**Contents**

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

**Immunogen**

A synthesized peptide derived from human PGK1

**Purification**

Affinity-chromatography

Storage

**Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.**

**Anti-PGK1 Rabbit Monoclonal Antibody - Protein Information**

**Name** PGK1

**Synonyms** PGKA

**Function**

Catalyzes one of the two ATP producing reactions in the glycolytic pathway via the reversible conversion of 1,3- diphosphoglycerate to 3-phosphoglycerate (PubMed:<a href="http://www.uniprot.org/citations/30323285" target="\_blank">30323285</a>, PubMed:<a href="http://www.uniprot.org/citations/7391028" target="\_blank">7391028</a>). In addition to its role as a glycolytic enzyme, it seems that PGK-1 acts as a polymerase alpha cofactor protein (primer recognition protein) (PubMed:<a href="http://www.uniprot.org/citations/2324090" target="\_blank">2324090</a>). May play a role in sperm motility (PubMed:<a href="http://www.uniprot.org/citations/26677959" target="\_blank">26677959</a>).

**Cellular Location**

Cytoplasm.

**Tissue Location**


Mainly expressed in spermatogonia. Localized on the principle piece in the sperm (at protein level). Expression significantly decreased in the testis of elderly men

**Anti-PGK1 Rabbit Monoclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Anti-PGK1 Rabbit Monoclonal Antibody - Images**



Western blot analysis of PGK1 expression in (1) HepG2 cell lysate; (2) Mouse kidney lysate.