

Anti-GFAP Rabbit Monoclonal Antibody
Catalog # ABO13387

Specification

Anti-GFAP Rabbit Monoclonal Antibody - Product Information

Application	WB, IHC, IF, ICC
Primary Accession	P14136
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human
Clonality	Monoclonal
Format	Liquid

Description

Anti-GFAP Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF applications. This antibody reacts with Human, Rat.

Anti-GFAP Rabbit Monoclonal Antibody - Additional Information

Gene ID 2670

Other Names

Glial fibrillary acidic protein, GFAP, GFAP

Calculated MW

49880 MW KDa

Application Details

WB 1:500-1:2000
IHC 1:50-1:200
ICC/IF 1:50-1:200

Subcellular Localization

Cytoplasm. Associated with intermediate filaments.

Tissue Specificity

Expressed in cells lacking fibronectin..

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human GFAP

Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Anti-GFAP Rabbit Monoclonal Antibody - Protein Information

Name GFAP

Function

GFAP, a class-III intermediate filament, is a cell-specific marker that, during the development of the central nervous system, distinguishes astrocytes from other glial cells.

Cellular Location

Cytoplasm. Note=Associated with intermediate filaments

Tissue Location


Expressed in cells lacking fibronectin.

Anti-GFAP Rabbit Monoclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


Anti-GFAP Rabbit Monoclonal Antibody - Images




All lanes use the Antibody at 1:2K dilution for 1 hour at room temperature.




All lanes use the Antibody at 1:2K dilution for 1 hour at room temperature.




All lanes use the Antibody at 1:2K dilution for 1 hour at room temperature.



Western blot analysis of GFAP expression in Rat brain lysate.



Immunohistochemical analysis of paraffin-embedded human glioma, using GFAP Antibody.



Immunofluorescent analysis of SNB19 cells, using GFAP Antibody.