

**Anti-PKM2 Rabbit Monoclonal Antibody**  
Catalog # ABO13433**Specification****Anti-PKM2 Rabbit Monoclonal Antibody - Product Information**

Application	WB, IHC, IF, ICC, FC
Primary Accession	<a href="#">P14618</a>
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

**Description**

Anti-PKM2 Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

**Anti-PKM2 Rabbit Monoclonal Antibody - Additional Information**

**Gene ID** 5315

**Other Names**

Pyruvate kinase PKM, 2.7.1.40, Cytosolic thyroid hormone-binding protein, CTHBP, Opa-interacting protein 3, OIP-3, Pyruvate kinase 2/3, Pyruvate kinase muscle isozyme, Threonine-protein kinase PKM2, 2.7.11.1, Thyroid hormone-binding protein 1, THBP1, Tumor M2-PK, Tyrosine-protein kinase PKM2, 2.7.10.2, p58, PKM, OIP3 {ECO:0000303|PubMed:9466265}, PK2, PK3, PKM2

**Calculated MW**

57937 MW KDa

**Application Details**

WB 1:1000-1:2000<br>IHC 1:50-1:200<br>ICC/IF 1:50-1:200<br>FC 1:50

**Subcellular Localization**

Cytoplasm. Nucleus. Translocates to the nucleus in response to different apoptotic stimuli. Nuclear translocation is sufficient to induce cell death that is caspase independent, isoform-specific and independent of its enzymatic activity.

**Tissue Specificity**

Specifically expressed in proliferating cells, such as embryonic stem cells, embryonic carcinoma cells, as well as cancer cells..

**Contents**

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

**Immunogen**

A synthesized peptide derived from human PKM2

**Purification**

Affinity-chromatography

Storage

**Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.**

## Anti-PKM2 Rabbit Monoclonal Antibody - Protein Information

**Name** PKM

**Synonyms** OIP3 {ECO:0000303|PubMed:9466265}, PK2,

### Function

Catalyzes the final rate-limiting step of glycolysis by mediating the transfer of a phosphoryl group from phosphoenolpyruvate (PEP) to ADP, generating ATP (PubMed:<a href="http://www.uniprot.org/citations/15996096" target="\_blank">15996096</a>, PubMed:<a href="http://www.uniprot.org/citations/1854723" target="\_blank">1854723</a>, PubMed:<a href="http://www.uniprot.org/citations/20847263" target="\_blank">20847263</a>). The ratio between the highly active tetrameric form and nearly inactive dimeric form determines whether glucose carbons are channeled to biosynthetic processes or used for glycolytic ATP production (PubMed:<a href="http://www.uniprot.org/citations/15996096" target="\_blank">15996096</a>, PubMed:<a href="http://www.uniprot.org/citations/1854723" target="\_blank">1854723</a>, PubMed:<a href="http://www.uniprot.org/citations/20847263" target="\_blank">20847263</a>). The transition between the 2 forms contributes to the control of glycolysis and is important for tumor cell proliferation and survival (PubMed:<a href="http://www.uniprot.org/citations/15996096" target="\_blank">15996096</a>, PubMed:<a href="http://www.uniprot.org/citations/1854723" target="\_blank">1854723</a>, PubMed:<a href="http://www.uniprot.org/citations/20847263" target="\_blank">20847263</a>).

### Cellular Location

[Isoform M2]: Cytoplasm. Nucleus Note=Translocates to the nucleus in response to various signals, such as EGF receptor activation or apoptotic stimuli (PubMed:17308100, PubMed:22056988, PubMed:24120661). Nuclear translocation is promoted by acetylation by EP300 (PubMed:24120661). Deacetylation by SIRT6 promotes its nuclear export in a process dependent of XPO4, thereby suppressing its ability to activate transcription and promote tumorigenesis (PubMed:26787900).

### Tissue Location


[Isoform M2]: Specifically expressed in proliferating cells, such as embryonic stem cells, embryonic carcinoma cells, as well as cancer cells.

## Anti-PKM2 Rabbit Monoclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## Anti-PKM2 Rabbit Monoclonal Antibody - Images



Western blot analysis of PKM2 expression in HeLa cell lysate.



Immunohistochemical analysis of paraffin-embedded human cervix cancer, using PKM2 Antibody.