

Anti-MRP2 ABCC2 Rabbit Monoclonal Antibody
Catalog # ABO13557**Specification****Anti-MRP2 ABCC2 Rabbit Monoclonal Antibody - Product Information**

Application	WB, IF, ICC, FC
Primary Accession	Q92887
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid

Description

Anti-MRP2 ABCC2 Rabbit Monoclonal Antibody . Tested in WB, ICC/IF, Flow Cytometry applications. This antibody reacts with Human.

Anti-MRP2 ABCC2 Rabbit Monoclonal Antibody - Additional Information

Gene ID 1244

Other Names

ATP-binding cassette sub-family C member 2, 7.6.2.-, 7.6.2.2, 7.6.2.3, Canalicular multidrug resistance protein, Canalicular multispecific organic anion transporter 1, Multidrug resistance-associated protein 2, ABCC2 (HGNC:53)

Calculated MW

174207 MW KDa

Application Details

WB 1:500-1:2000
ICC/IF 1:50-1:200
FC 1:100

Subcellular Localization

Apical cell membrane ; Multi-pass membrane protein.

Tissue Specificity

Expressed by polarized cells in liver, kidney and intestine. The highest expression is found in liver.

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human MRP2

Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Anti-MRP2 ABCC2 Rabbit Monoclonal Antibody - Protein Information

Name ABCC2 ([HGNC:53](#))

Function

ATP-dependent transporter of the ATP-binding cassette (ABC) family that binds and hydrolyzes ATP to enable active transport of various substrates including many drugs, toxicants and endogenous compound across cell membranes. Transports a wide variety of conjugated organic anions such as sulfate-, glucuronide- and glutathione (GSH)- conjugates of endo- and xenobiotics substrates (PubMed:[10220572](http://www.uniprot.org/citations/10220572)), PubMed:[10421658](http://www.uniprot.org/citations/10421658)), PubMed:[11500505](http://www.uniprot.org/citations/11500505)), PubMed:[16332456](http://www.uniprot.org/citations/16332456)). Mediates hepatobiliary excretion of mono- and bis-glucuronidated bilirubin molecules and therefore play an important role in bilirubin detoxification (PubMed:[10421658](http://www.uniprot.org/citations/10421658)). Mediates also hepatobiliary excretion of others glucuronide conjugates such as 17beta-estradiol 17-glucosiduronic acid and leukotriene C4 (PubMed:[11500505](http://www.uniprot.org/citations/11500505)). Transports sulfated bile salt such as tauroolithocholate sulfate (PubMed:[16332456](http://www.uniprot.org/citations/16332456)). Transports various anticancer drugs, such as anthracycline, vinca alkaloid and methotrexate and HIV-drugs such as protease inhibitors (PubMed:[10220572](http://www.uniprot.org/citations/10220572)), PubMed:[11500505](http://www.uniprot.org/citations/11500505)), PubMed:[12441801](http://www.uniprot.org/citations/12441801)). Confers resistance to several anti-cancer drugs including cisplatin, doxorubicin, epirubicin, methotrexate, etoposide and vincristine (PubMed:[10220572](http://www.uniprot.org/citations/10220572)), PubMed:[11500505](http://www.uniprot.org/citations/11500505)).

Cellular Location

Apical cell membrane; Multi-pass membrane protein. Note=Localized to the apical membrane of enterocytes

Tissue Location

Expressed by polarized cells in liver, kidney and intestine. The highest expression is found in liver. Expressed in small intestine (PubMed:28408210).

Anti-MRP2 ABCC2 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)

• [Cell Culture](#)

Anti-MRP2 ABCC2 Rabbit Monoclonal Antibody - Images




Figure 1. Western blot analysis of ABCC2 using anti-ABCC2 antibody (M00974).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human HepG2 whole cell lysates,

Lane 2: human A549 whole cell lysates,

Lane 3: human CACO-2 whole cell lysates,

Lane 4: human HUH-7 whole cell lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-ABCC2 antigen affinity purified monoclonal antibody (Catalog # M00974) at 1:500 overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:500 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for ABCC2 at approximately 250 kDa. The expected band size for ABCC2 is at 174 kDa.