

## Anti-ROCK1 Rabbit Monoclonal Antibody Catalog # ABO13624

### Specification

---

#### Anti-ROCK1 Rabbit Monoclonal Antibody - Product Information

Application	WB, IHC, IF, ICC, IP, FC
Primary Accession	<a href="#">Q13464</a>
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

#### Description

Anti-ROCK1 Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

#### Anti-ROCK1 Rabbit Monoclonal Antibody - Additional Information

**Gene ID** 6093

#### Other Names

Rho-associated protein kinase 1, 2.7.11.1, Renal carcinoma antigen NY-REN-35, Rho-associated, coiled-coil-containing protein kinase 1, Rho-associated, coiled-coil-containing protein kinase I, ROCK-I, p160 ROCK-1, p160ROCK, ROCK1

#### Calculated MW

158175 MW KDa

#### Application Details

WB 1:500-1:2000<br>IHC 1:50-1:200<br>ICC/IF 1:50-1:200<br>IP 1:50<br>FC 1:50

#### Subcellular Localization

Cytoplasm. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome, centriole. Golgi apparatus membrane; Peripheral membrane protein. Cell projection, bleb. Cytoplasm, cytoskeleton. Cell membrane. Cell projection, lamellipodium. Cell projection, ruffle. Associated with the mother centriole and an intercentriolar linker. Colocalizes with ITGB1BP1 and ITGB1 at the cell membrane predominantly in lamellipodia and membrane ruffles, but also in retraction fibers. Localizes at the cell membrane in an ITGB1BP1-dependent manner (By similarity). A small proportion is associated with Golgi membranes..

#### Tissue Specificity

Detected in blood platelets..

#### Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

#### Immunogen

A synthesized peptide derived from human ROCK1

## Purification

Affinity-chromatography

Storage

**Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.**

## Anti-ROCK1 Rabbit Monoclonal Antibody - Protein Information

Name ROCK1

### Function

Protein kinase which is a key regulator of the actin cytoskeleton and cell polarity (PubMed:<a href="http://www.uniprot.org/citations/10436159" target="\_blank">10436159</a>, PubMed:<a href="http://www.uniprot.org/citations/10652353" target="\_blank">10652353</a>, PubMed:<a href="http://www.uniprot.org/citations/11018042" target="\_blank">11018042</a>, PubMed:<a href="http://www.uniprot.org/citations/11283607" target="\_blank">11283607</a>, PubMed:<a href="http://www.uniprot.org/citations/17158456" target="\_blank">17158456</a>, PubMed:<a href="http://www.uniprot.org/citations/18573880" target="\_blank">18573880</a>, PubMed:<a href="http://www.uniprot.org/citations/19131646" target="\_blank">19131646</a>, PubMed:<a href="http://www.uniprot.org/citations/8617235" target="\_blank">8617235</a>, PubMed:<a href="http://www.uniprot.org/citations/9722579" target="\_blank">9722579</a>). Involved in regulation of smooth muscle contraction, actin cytoskeleton organization, stress fiber and focal adhesion formation, neurite retraction, cell adhesion and motility via phosphorylation of DAPK3, GFAP, LIMK1, LIMK2, MYL9/MLC2, TPPP, PFN1 and PPP1R12A (PubMed:<a href="http://www.uniprot.org/citations/10436159" target="\_blank">10436159</a>, PubMed:<a href="http://www.uniprot.org/citations/10652353" target="\_blank">10652353</a>, PubMed:<a href="http://www.uniprot.org/citations/11018042" target="\_blank">11018042</a>, PubMed:<a href="http://www.uniprot.org/citations/11283607" target="\_blank">11283607</a>, PubMed:<a href="http://www.uniprot.org/citations/17158456" target="\_blank">17158456</a>, PubMed:<a href="http://www.uniprot.org/citations/18573880" target="\_blank">18573880</a>, PubMed:<a href="http://www.uniprot.org/citations/19131646" target="\_blank">19131646</a>, PubMed:<a href="http://www.uniprot.org/citations/23093407" target="\_blank">23093407</a>, PubMed:<a href="http://www.uniprot.org/citations/23355470" target="\_blank">23355470</a>, PubMed:<a href="http://www.uniprot.org/citations/8617235" target="\_blank">8617235</a>, PubMed:<a href="http://www.uniprot.org/citations/9722579" target="\_blank">9722579</a>). Phosphorylates FHOD1 and acts synergistically with it to promote SRC-dependent non-apoptotic plasma membrane blebbing (PubMed:<a href="http://www.uniprot.org/citations/18694941" target="\_blank">18694941</a>). Phosphorylates JIP3 and regulates the recruitment of JNK to JIP3 upon UVB-induced stress (PubMed:<a href="http://www.uniprot.org/citations/19036714" target="\_blank">19036714</a>). Acts as a suppressor of inflammatory cell migration by regulating PTEN phosphorylation and stability (By similarity). Acts as a negative regulator of VEGF-induced angiogenic endothelial cell activation (PubMed:<a href="http://www.uniprot.org/citations/19181962" target="\_blank">19181962</a>). Required for centrosome positioning and centrosome-dependent exit from mitosis (By similarity). Plays a role in terminal erythroid differentiation (PubMed:<a href="http://www.uniprot.org/citations/21072057" target="\_blank">21072057</a>). Inhibits podocyte motility via regulation of actin cytoskeletal dynamics and phosphorylation of CFL1 (By similarity). Promotes keratinocyte terminal differentiation (PubMed:<a href="http://www.uniprot.org/citations/19997641" target="\_blank">19997641</a>). Involved in osteoblast compaction through the fibronectin fibrillogenesis cell-mediated matrix assembly process, essential for osteoblast mineralization (By similarity). May regulate closure of the eyelids and ventral body wall by inducing the assembly of actomyosin bundles (By similarity).

### Cellular Location

Cytoplasm. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome, centriole {ECO:0000250|UniProtKB:P70335}. Golgi apparatus membrane; Peripheral membrane protein. Cell projection, bleb. Cytoplasm, cytoskeleton {ECO:0000250|UniProtKB:P70335}. Cell membrane {ECO:0000250|UniProtKB:P70335}. Cell projection, lamellipodium {ECO:0000250|UniProtKB:P70335}. Cell projection, ruffle {ECO:0000250|UniProtKB:P70335}. Note=A small proportion is associated with Golgi membranes (PubMed:12773565). Associated with the mother centriole and an intercentriolar linker (By similarity). Colocalizes with ITGB1BP1 and ITGB1 at the cell membrane predominantly in lamellipodia and membrane ruffles, but also in retraction fibers (By similarity). Localizes at the cell membrane in an ITGB1BP1-dependent manner (By similarity). {ECO:0000250|UniProtKB:P70335, ECO:0000269|PubMed:12773565}

#### Tissue Location

Detected in blood platelets.

### Anti-ROCK1 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Anti-ROCK1 Rabbit Monoclonal Antibody - Images





Figure 1. Western blot analysis of ROCK1 using anti-ROCK1 antibody (M00722).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.


Lane 1: human HeLa whole cell lysates,  
Lane 2: human Jurkat whole cell lysates,  
Lane 3: human U2OS whole cell lysates,  
Lane 4: human 293T whole cell lysates,  
Lane 5: rat brain tissue lysates,

Lane 6: rat C6 whole cell lysates,  
Lane 7: mouse brain tissue lysates,  
Lane 8: mouse Neuro-2a whole cell lysates.


After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-ROCK1 antigen affinity purified monoclonal antibody (Catalog # M00722) at 1:500 overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:500 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for ROCK1 at approximately 170 kDa. The expected band size for ROCK1 is at 158 kDa.




Immunohistochemical analysis of paraffin-embedded Rat stomach, using the Antibody at 1:150 dilution.




Immunohistochemical analysis of paraffin-embedded Rat heart, using the Antibody at 1:150 dilution.




Immunohistochemical analysis of paraffin-embedded Human squamous carcinoma, using the Antibody at 1:150 dilution.




Immunohistochemical analysis of paraffin-embedded Human esophageal carcinoma, using the Antibody at 1:150 dilution.



Immunohistochemical analysis of paraffin-embedded Mouse heart, using the Antibody at 1:150 dilution.



Immunofluorescent analysis of HeLa cells, using ROCK1 Antibody .



Immunofluorescent analysis using the Antibody at 1:50 dilution.