

Anti-VCAM1/Cd106 Rabbit Monoclonal Antibody
Catalog # ABO13637

Specification

Anti-VCAM1/Cd106 Rabbit Monoclonal Antibody - Product Information

Application	WB, IHC, IF, ICC, IP, FC
Primary Accession	P19320
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

Description

Anti-VCAM1/Cd106 Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

Anti-VCAM1/Cd106 Rabbit Monoclonal Antibody - Additional Information

Gene ID 7412

Other Names

Vascular cell adhesion protein 1, V-CAM 1, VCAM-1, INCAM-100, CD106, Soluble Vascular Cell Adhesion Molecule-1, VCAM1

Calculated MW

81276 MW KDa

Application Details

WB 1:500-1:2000
IHC 1:50-1:200
ICC/IF 1:50-1:200
IP 1:20
FC 1:50

Subcellular Localization

Membrane; Single-pass type I membrane protein.

Tissue Specificity

Expressed on inflammed vascular endothelium, as well as on macrophage-like and dendritic cell types in both normal and inflammed tissue.

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human VCAM1

Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for

up to one month. Avoid repeated
freeze-thaw cycles.

Anti-VCAM1/Cd106 Rabbit Monoclonal Antibody - Protein Information

Name VCAM1

Function

Cell adhesion glycoprotein predominantly expressed on the surface of endothelial cells that plays an important role in immune surveillance and inflammation (PubMed:31310649). Acts as a major regulator of leukocyte adhesion to the endothelium through interaction with different types of integrins (PubMed:10209034). During inflammatory responses, binds ligands on the surface of activated endothelial cells to initiate the activation of calcium channels and the plasma membrane-associated small GTPase RAC1 leading to leukocyte transendothelial migration (PubMed:22970700). Serves also as a quality- control checkpoint for entry into bone marrow by providing a 'don't-eat-me' stamping in the context of major histocompatibility complex (MHC) class-I presentation (PubMed:35210567).

Cellular Location

[Vascular cell adhesion protein 1]: Cell membrane; Single-pass type I membrane protein

Tissue Location

Expressed on inflamed vascular endothelium, as well as on macrophage-like and dendritic cell types in both normal and inflamed tissue


Anti-VCAM1/Cd106 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-VCAM1/Cd106 Rabbit Monoclonal Antibody - Images






Western blot analysis of VCAM1 expression in Mouse kidney lysate.



Immunohistochemical analysis of paraffin-embedded human tonsil, using VCAM1 Antibody.



Immunofluorescent analysis of K562 cells, using VCAM1 Antibody.