

Anti-SOX18 Rabbit Monoclonal Antibody

Catalog # ABO13638

Specification

Anti-SOX18 Rabbit Monoclonal Antibody - Product Information

Application	WB. IHC	
Primary Accession	P35713	
Host	Rabbit	
Isotype	Rabbit IgG	
Reactivity	Human	
Clonality	Monoclonal	
Format	Liquid	
Description	•	
Anti-SOX18 Rabbit Monoclonal Antibo	dy. Tested in WB, IHC applications. This antibody reacts wit	th
Human.		

Anti-SOX18 Rabbit Monoclonal Antibody - Additional Information

Gene ID 54345

Other Names Transcription factor SOX-18, SOX18

Calculated MW 40891 MW KDa

Application Details WB 1:500-1:2000
HC 1:50-1:200

Subcellular Localization Nucleus.

Contents Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen A synthesized peptide derived from human SOX18

Purification Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Anti-SOX18 Rabbit Monoclonal Antibody - Protein Information



Name SOX18

Function

Transcriptional activator that binds to the consensus sequence 5'-AACAAAG-3' in the promoter of target genes and plays an essential role in embryonic cardiovascular development and lymphangiogenesis. Activates transcription of PROX1 and other genes coding for lymphatic endothelial markers. Plays an essential role in triggering the differentiation of lymph vessels, but is not required for the maintenance of differentiated lymphatic endothelial cells. Plays an important role in postnatal angiogenesis, where it is functionally redundant with SOX17. Interaction with MEF2C enhances transcriptional activation. Besides, required for normal hair development.

Cellular Location Nucleus {ECO:0000255|PROSITE-ProRule:PRU00267}.

Tissue Location

Detected in heart, lung, placenta, skeletal muscle, liver, kidney, spleen, prostate, ovary, msosmall intestine and colon

Anti-SOX18 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Anti-SOX18 Rabbit Monoclonal Antibody - Images



Figure 1. Western blot analysis of SOX18 using anti-SOX18 antibody (M04004).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human K562 whole cell lysates,



Lane 2: human COLO 320 whole cell lysates,

Lane 3: human HepG2 whole cell lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-SOX18 antigen affinity purified monoclonal antibody (Catalog # M04004) at 1:500 overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for SOX18 at approximately 41 kDa.