

**Anti-LAMP1/Cd107A Rabbit Monoclonal Antibody**  
Catalog # ABO13727**Specification****Anti-LAMP1/Cd107A Rabbit Monoclonal Antibody - Product Information**

Application	WB, IHC
Primary Accession	<a href="#">P11279</a>
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid

**Description**

Anti-LAMP1/Cd107A Rabbit Monoclonal Antibody . Tested in WB, IHC applications. This antibody reacts with Human.

**Anti-LAMP1/Cd107A Rabbit Monoclonal Antibody - Additional Information**

**Gene ID** 3916

**Other Names**

Lysosome-associated membrane glycoprotein 1, LAMP-1, Lysosome-associated membrane protein 1, CD107 antigen-like family member A, CD107a, LAMP1 {ECO:0000303|PubMed:23632890, ECO:0000312|HGNC:HGNC:6499}

**Calculated MW**

44882 MW KDa

**Application Details**

WB 1:500-1:2000<br>IHC 1:50-1:200

**Subcellular Localization**

Cell membrane; Single-pass type I membrane protein. Endosome membrane; Single-pass type I membrane protein. Lysosome membrane; Single-pass type I membrane protein. Late endosome. This protein shuttles between lysosomes, endosomes, and the plasma membrane. Colocalizes with OSBPL1A at the late endosome.

**Contents**

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

**Immunogen**

A synthesized peptide derived from human LAMP1

**Purification**

Affinity-chromatography

Storage

**Store at -20°C for one year. For short term storage and frequent use, store at 4°C for**

**up to one month. Avoid repeated  
freeze-thaw cycles.**

## **Anti-LAMP1/Cd107A Rabbit Monoclonal Antibody - Protein Information**

**Name** LAMP1 {ECO:0000303|PubMed:23632890, ECO:0000312|HGNC:HGNC:6499}

### **Function**

Lysosomal membrane glycoprotein which plays an important role in lysosome biogenesis, lysosomal pH regulation, autophagy and cholesterol homeostasis (PubMed:<a href="http://www.uniprot.org/citations/37390818" target="\_blank">37390818</a>). Acts as an important regulator of lysosomal lumen pH regulation by acting as a direct inhibitor of the proton channel TMEM175, facilitating lysosomal acidification for optimal hydrolase activity (PubMed:<a href="http://www.uniprot.org/citations/37390818" target="\_blank">37390818</a>). Also plays an important role in NK-cells cytotoxicity (PubMed:<a href="http://www.uniprot.org/citations/2022921" target="\_blank">2022921</a>, PubMed:<a href="http://www.uniprot.org/citations/23632890" target="\_blank">23632890</a>). Mechanistically, participates in cytotoxic granule movement to the cell surface and perforin trafficking to the lytic granule (PubMed:<a href="http://www.uniprot.org/citations/23632890" target="\_blank">23632890</a>). In addition, protects NK-cells from degranulation-associated damage induced by their own cytotoxic granule content (PubMed:<a href="http://www.uniprot.org/citations/23847195" target="\_blank">23847195</a>). Presents carbohydrate ligands to selectins (PubMed:<a href="http://www.uniprot.org/citations/7685349" target="\_blank">7685349</a>).

### **Cellular Location**


Lysosome membrane; Single-pass type I membrane protein. Endosome membrane; Single-pass type I membrane protein. Late endosome membrane; Single-pass type I membrane protein. Cell membrane; Single-pass type I membrane protein. Cytolytic granule membrane; Single-pass type I membrane protein. Note=This protein shuttles between lysosomes, endosomes, and the plasma membrane (By similarity). Colocalizes with OSBPL1A at the late endosome (PubMed:16176980). {ECO:0000250|UniProtKB:P05300, ECO:0000269|PubMed:16176980, ECO:0000269|PubMed:17897319}


## **Anti-LAMP1/Cd107A Rabbit Monoclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## **Anti-LAMP1/Cd107A Rabbit Monoclonal Antibody - Images**





All lanes use the Antibody at 1:6K dilution for 1 hour at room temperature.



Western blot analysis of LAMP1 expression in A431 cell lysate.