

Anti-PRDX6/Peroxiredoxin 6 Rabbit Monoclonal Antibody
Catalog # ABO13774

Specification

Anti-PRDX6/Peroxiredoxin 6 Rabbit Monoclonal Antibody - Product Information

Application	WB, IF, ICC, IP
Primary Accession	P30041
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

Description

Anti-PRDX6/Peroxiredoxin 6 Rabbit Monoclonal Antibody . Tested in WB, ICC/IF, IP applications. This antibody reacts with Human, Mouse, Rat.

Anti-PRDX6/Peroxiredoxin 6 Rabbit Monoclonal Antibody - Additional Information

Gene ID 9588

Other Names

Peroxiredoxin-6, 1.11.1.27, 1-Cys peroxiredoxin, 1-Cys PRX, 24 kDa protein, Acidic calcium-independent phospholipase A2, aiPLA2, 3.1.1.4, Antioxidant protein 2, Glutathione-dependent peroxiredoxin, Liver 2D page spot 40, Lysophosphatidylcholine acyltransferase 5, LPC acyltransferase 5, LPCAT-5, Lyso-PC acyltransferase 5, 2.3.1.23, Non-selenium glutathione peroxidase, NSGPx, Red blood cells page spot 12, PRDX6, AOP2, KIAA0106

Calculated MW

25035 MW KDa

Application Details

WB 1:500-1:1000
ICC/IF 1:50-1:200
IP 1:20

Subcellular Localization

Cytoplasm. Lysosome. Cytoplasmic vesicle. Also found in lung secretory organelles..

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human PRDX6

Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for

up to one month. Avoid repeated
freeze-thaw cycles.

Anti-PRDX6/Peroxiredoxin 6 Rabbit Monoclonal Antibody - Protein Information

Name PRDX6

Synonyms AOP2, KIAA0106

Function

Thiol-specific peroxidase that catalyzes the reduction of hydrogen peroxide and organic hydroperoxides to water and alcohols, respectively (PubMed: [10893423](http://www.uniprot.org/citations/10893423), PubMed: [9497358](http://www.uniprot.org/citations/9497358)). Can reduce H₂O₂ and short chain organic, fatty acid, and phospholipid hydroperoxides (PubMed: [10893423](http://www.uniprot.org/citations/10893423)). Also has phospholipase activity, can therefore either reduce the oxidized sn-2 fatty acyl group of phospholipids (peroxidase activity) or hydrolyze the sn-2 ester bond of phospholipids (phospholipase activity) (PubMed: [10893423](http://www.uniprot.org/citations/10893423), PubMed: [26830860](http://www.uniprot.org/citations/26830860)). These activities are dependent on binding to phospholipids at acidic pH and to oxidized phospholipids at cytosolic pH (PubMed: [10893423](http://www.uniprot.org/citations/10893423)). Plays a role in cell protection against oxidative stress by detoxifying peroxides and in phospholipid homeostasis (PubMed: [10893423](http://www.uniprot.org/citations/10893423)). Exhibits acyl-CoA-dependent lysophospholipid acyltransferase which mediates the conversion of lysophosphatidylcholine (1-acyl-sn-glycero-3-phosphocholine or LPC) into phosphatidylcholine (1,2-diacyl-sn-glycero-3-phosphocholine or PC) (PubMed: [26830860](http://www.uniprot.org/citations/26830860)). Shows a clear preference for LPC as the lysophospholipid and for palmitoyl CoA as the fatty acyl substrate (PubMed: [26830860](http://www.uniprot.org/citations/26830860)).

Cellular Location


Cytoplasm. Lysosome {ECO:0000250|UniProtKB:O35244}. Note=Also found in lung secretory organelles (lamellar bodies). {ECO:0000250|UniProtKB:O35244}

Anti-PRDX6/Peroxiredoxin 6 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-PRDX6/Peroxiredoxin 6 Rabbit Monoclonal Antibody - Images






Figure 1. Western blot analysis of PRDX6 using anti-PRDX6 antibody (M01847).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

- Lane 1: human PC-3 whole cell lysates,
- Lane 2: human K562 whole cell lysates,
- Lane 3: human Hela whole cell lysates,
- Lane 4: rat brain tissue lysates,
- Lane 5: rat liver tissue lysates,
- Lane 6: mouse brain tissue lysates,
- Lane 7: mouse liver tissue lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-PRDX6 antigen affinity purified monoclonal antibody (Catalog # M01847) at 1:500 overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for PRDX6 at approximately 25 kDa. The expected band size for PRDX6 is at 25 kDa.