

## Anti-GATA4 Rabbit Monoclonal Antibody Catalog # ABO13781

### Specification

---

#### Anti-GATA4 Rabbit Monoclonal Antibody - Product Information

Application	WB, IF, ICC, FC
Primary Accession	<a href="#">P43694</a>
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

#### Description

Anti-GATA4 Rabbit Monoclonal Antibody . Tested in WB, ICC/IF, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

#### Anti-GATA4 Rabbit Monoclonal Antibody - Additional Information

**Gene ID** 2626

#### Other Names

Transcription factor GATA-4, GATA-binding factor 4, GATA4

#### Calculated MW

44565 MW KDa

#### Application Details

WB 1:500-1:2000<br>ICC/IF 1:50-1:200<br>FC 1:100

#### Subcellular Localization

Nucleus.

#### Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

#### Immunogen

A synthesized peptide derived from human GATA4

#### Purification

Affinity-chromatography

#### Storage

**Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.**

#### Anti-GATA4 Rabbit Monoclonal Antibody - Protein Information

## Name GATA4

### Function

Transcriptional activator that binds to the consensus sequence 5'-AGATAG-3' and plays a key role in cardiac development and function (PubMed:<a href="http://www.uniprot.org/citations/24000169" target="\_blank">24000169</a>, PubMed:<a href="http://www.uniprot.org/citations/27984724" target="\_blank">27984724</a>, PubMed:<a href="http://www.uniprot.org/citations/35182466" target="\_blank">35182466</a>). In cooperation with TBX5, it binds to cardiac super-enhancers and promotes cardiomyocyte gene expression, while it down-regulates endocardial and endothelial gene expression (PubMed:<a href="http://www.uniprot.org/citations/27984724" target="\_blank">27984724</a>). Involved in bone morphogenetic protein (BMP)-mediated induction of cardiac-specific gene expression. Binds to BMP response element (BMPRE) DNA sequences within cardiac activating regions (By similarity). Acts as a transcriptional activator of ANF in cooperation with NKX2-5 (By similarity). Promotes cardiac myocyte enlargement (PubMed:<a href="http://www.uniprot.org/citations/20081228" target="\_blank">20081228</a>). Required during testicular development (PubMed:<a href="http://www.uniprot.org/citations/21220346" target="\_blank">21220346</a>). May play a role in sphingolipid signaling by regulating the expression of sphingosine-1-phosphate degrading enzyme, sphingosine-1-phosphate lyase (PubMed:<a href="http://www.uniprot.org/citations/15734735" target="\_blank">15734735</a>).

### Cellular Location


Nucleus

## Anti-GATA4 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## Anti-GATA4 Rabbit Monoclonal Antibody - Images



Immunofluorescent analysis of HepG2 cells, using GATA4 Antibody .




Figure 1. Western blot analysis of GATA4 using anti-GATA4 antibody (M00499-1).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human HepG2 whole cell lysates,

Lane 2: human SH-SY5Y whole cell lysates,

Lane 3: rat liver tissue lysates,

Lane 4: mouse liver tissue lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-GATA4 antigen affinity purified monoclonal antibody (Catalog # M00499-1) at 1:500 overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for GATA4 at approximately 54 kDa. The expected band size for GATA4 is at 45 kDa.