

Anti-CD276/B7 H3 Rabbit Monoclonal Antibody
Catalog # ABO13801**Specification****Anti-CD276/B7 H3 Rabbit Monoclonal Antibody - Product Information**

Application	WB, IP, FC
Primary Accession	Q5ZPR3
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Human, Mouse
Clonality	Monoclonal
Format	Liquid

Description

Anti-CD276/B7 H3 Rabbit Monoclonal Antibody . Tested in WB, IP, Flow Cytometry applications. This antibody reacts with Human, Mouse.

Anti-CD276/B7 H3 Rabbit Monoclonal Antibody - Additional Information

Gene ID 80381

Other Names

CD276 antigen, 4lg-B7-H3, B7 homolog 3, B7-H3, Costimulatory molecule, CD276, CD276, B7H3

Calculated MW

57235 MW KDa

Application Details

WB 1:500-1:2000
IP 1:50
FC 1:200

Subcellular Localization

Membrane ; Single-pass type I membrane protein.

Tissue Specificity

Ubiquitous but not detectable in peripheral blood lymphocytes or granulocytes. Weakly expressed in resting monocytes. Expressed in dendritic cells derived from monocytes. Expressed in epithelial cells of sinonasal tissue. Expressed in extravillous trophoblast cells and Hofbauer cells of the first trimester placenta and term placenta..

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human CD276

Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term

storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Anti-CD276/B7 H3 Rabbit Monoclonal Antibody - Protein Information

Name CD276

Synonyms B7H3

Function

May participate in the regulation of T-cell-mediated immune response. May play a protective role in tumor cells by inhibiting natural-killer mediated cell lysis as well as a role of marker for detection of neuroblastoma cells. May be involved in the development of acute and chronic transplant rejection and in the regulation of lymphocytic activity at mucosal surfaces. Could also play a key role in providing the placenta and fetus with a suitable immunological environment throughout pregnancy. Both isoform 1 and isoform 2 appear to be redundant in their ability to modulate CD4 T-cell responses. Isoform 2 is shown to enhance the induction of cytotoxic T-cells and selectively stimulates interferon gamma production in the presence of T-cell receptor signaling.

Cellular Location

Membrane; Single-pass type I membrane protein

Tissue Location


Ubiquitous but not detectable in peripheral blood lymphocytes or granulocytes. Weakly expressed in resting monocytes Expressed in dendritic cells derived from monocytes. Expressed in epithelial cells of sinonasal tissue. Expressed in extravillous trophoblast cells and Hofbauer cells of the first trimester placenta and term placenta.


Anti-CD276/B7 H3 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-CD276/B7 H3 Rabbit Monoclonal Antibody - Images





Western blot analysis of CD276 expression in 293T cell lysate.