

Anti-CYP2E1 Rabbit Monoclonal Antibody
Catalog # ABO13870

Specification

Anti-CYP2E1 Rabbit Monoclonal Antibody - Product Information

Application	WB
Primary Accession	P05181
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

Description

Anti-CYP2E1 Rabbit Monoclonal Antibody . Tested in WB application. This antibody reacts with Human, Mouse, Rat.

Anti-CYP2E1 Rabbit Monoclonal Antibody - Additional Information

Gene ID 1571

Other Names

Cytochrome P450 2E1, 1.14.14.1, 4-nitrophenol 2-hydroxylase, 1.14.13.n7, CYP11E1, Cytochrome P450-J, CYP2E1 {ECO:0000303|PubMed:10553002, ECO:0000312|HGNC:HGNC:2631}

Calculated MW

56849 MW KDa

Application Details

WB 1:500-1:2000

Subcellular Localization

Endoplasmic reticulum membrane; Peripheral membrane protein. Microsome membrane; Peripheral membrane protein.

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human CYP2E1

Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Anti-CYP2E1 Rabbit Monoclonal Antibody - Protein Information

Name CYP2E1 {ECO:0000303|PubMed:10553002, ECO:0000312|HGNC:HGNC:2631}

Function

A cytochrome P450 monooxygenase involved in the metabolism of fatty acids (PubMed:10553002, PubMed:18577768). Mechanistically, uses molecular oxygen inserting one oxygen atom into a substrate, and reducing the second into a water molecule, with two electrons provided by NADPH via cytochrome P450 reductase (NADPH--hemoprotein reductase) (PubMed:10553002, PubMed:18577768). Catalyzes the hydroxylation of carbon-hydrogen bonds. Hydroxylates fatty acids specifically at the omega-1 position displaying the highest catalytic activity for saturated fatty acids (PubMed:10553002, PubMed:18577768). May be involved in the oxidative metabolism of xenobiotics (Probable).

Cellular Location


Endoplasmic reticulum membrane {ECO:0000250|UniProtKB:P05182}; Peripheral membrane protein {ECO:0000250|UniProtKB:P05182}. Microsome membrane {ECO:0000250|UniProtKB:P05182}; Peripheral membrane protein {ECO:0000250|UniProtKB:P05182}. Mitochondrion inner membrane {ECO:0000250|UniProtKB:P05182}; Peripheral membrane protein {ECO:0000250|UniProtKB:P05182}. Note=Post-translationally targeted to mitochondria. TOMM70 is required for the translocation across the mitochondrial outer membrane. After translocation into the matrix, associates with the inner membrane as a membrane extrinsic protein {ECO:0000250|UniProtKB:P05182}


Anti-CYP2E1 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-CYP2E1 Rabbit Monoclonal Antibody - Images





Western blot analysis of CYP2E1 expression in HeLa cell lysate.