

**Anti-HLA-DRA/Hla Dr Rabbit Monoclonal Antibody**  
Catalog # ABO13897**Specification****Anti-HLA-DRA/Hla Dr Rabbit Monoclonal Antibody - Product Information**

Application	WB, IHC, IF, ICC, FC
Primary Accession	<a href="#">P01903</a>
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid

**Description**

Anti-HLA-DRA/Hla Dr Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, Flow Cytometry applications. This antibody reacts with Human.

**Anti-HLA-DRA/Hla Dr Rabbit Monoclonal Antibody - Additional Information**

**Gene ID** 3122

**Other Names**

HLA class II histocompatibility antigen, DR alpha chain, MHC class II antigen DRA, HLA-DRA, HLA-DRA1

**Calculated MW**

28607 MW KDa

**Application Details**

WB 1:1000-1:5000<br>IHC 1:50-1:200<br>ICC/IF 1:50-1:200<br>FC 1:50

**Subcellular Localization**

Cell membrane; Single-pass type I membrane protein. Endoplasmic reticulum membrane; Single-pass type I membrane protein. Golgi apparatus, trans-Golgi network membrane; Single-pass type I membrane protein. Endosome membrane; Single-pass type I membrane protein. Lysosome membrane; Single-pass type I membrane protein. Late endosome membrane; Single-pass type I membrane protein. The MHC class II complex transits through a number of intracellular compartments in the endocytic pathway until it reaches the cell membrane for antigen presentation.

**Contents**

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

**Immunogen**

A synthesized peptide derived from human HLA-DRA

**Purification**

Affinity-chromatography

**Storage**

**Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.**

**Anti-HLA-DRA/Hla Dr Rabbit Monoclonal Antibody - Protein Information****Name** HLA-DRA**Synonyms** HLA-DRA1**Function**

An alpha chain of antigen-presenting major histocompatibility complex class II (MHCII) molecule. In complex with the beta chain HLA- DRB, displays antigenic peptides on professional antigen presenting cells (APCs) for recognition by alpha-beta T cell receptor (TCR) on HLA-DR-restricted CD4-positive T cells. This guides antigen-specific T- helper effector functions, both antibody-mediated immune response and macrophage activation, to ultimately eliminate the infectious agents and transformed cells (PubMed:<a href="http://www.uniprot.org/citations/15265931" target="\_blank">15265931</a>, PubMed:<a href="http://www.uniprot.org/citations/15322540" target="\_blank">15322540</a>, PubMed:<a href="http://www.uniprot.org/citations/17334368" target="\_blank">17334368</a>, PubMed:<a href="http://www.uniprot.org/citations/22327072" target="\_blank">22327072</a>, PubMed:<a href="http://www.uniprot.org/citations/24190431" target="\_blank">24190431</a>, PubMed:<a href="http://www.uniprot.org/citations/27591323" target="\_blank">27591323</a>, PubMed:<a href="http://www.uniprot.org/citations/29884618" target="\_blank">29884618</a>, PubMed:<a href="http://www.uniprot.org/citations/31495665" target="\_blank">31495665</a>, PubMed:<a href="http://www.uniprot.org/citations/8145819" target="\_blank">8145819</a>, PubMed:<a href="http://www.uniprot.org/citations/9075930" target="\_blank">9075930</a>). Typically presents extracellular peptide antigens of 10 to 30 amino acids that arise from proteolysis of endocytosed antigens in lysosomes (PubMed:<a href="http://www.uniprot.org/citations/8145819" target="\_blank">8145819</a>). In the tumor microenvironment, presents antigenic peptides that are primarily generated in tumor-resident APCs likely via phagocytosis of apoptotic tumor cells or macropinocytosis of secreted tumor proteins (PubMed:<a href="http://www.uniprot.org/citations/31495665" target="\_blank">31495665</a>). Presents peptides derived from intracellular proteins that are trapped in autolysosomes after macroautophagy, a mechanism especially relevant for T cell selection in the thymus and central immune tolerance (PubMed:<a href="http://www.uniprot.org/citations/17182262" target="\_blank">17182262</a>, PubMed:<a href="http://www.uniprot.org/citations/23783831" target="\_blank">23783831</a>). The selection of the immunodominant epitopes follows two processing modes: 'bind first, cut/trim later' for pathogen-derived antigenic peptides and 'cut first, bind later' for autoantigens/self- peptides (PubMed:<a href="http://www.uniprot.org/citations/25413013" target="\_blank">25413013</a>). The anchor residue at position 1 of the peptide N-terminus, usually a large hydrophobic residue, is essential for high affinity interaction with MHCII molecules (PubMed:<a href="http://www.uniprot.org/citations/8145819" target="\_blank">8145819</a>).

**Cellular Location**

Cell membrane; Single-pass type I membrane protein. Endoplasmic reticulum membrane; Single-pass type I membrane protein. Early endosome membrane; Single-pass type I membrane protein. Late endosome membrane; Single-pass type I membrane protein. Lysosome membrane; Single-pass type I membrane protein. Autolysosome membrane; Single-pass type I membrane protein. Note=The MHCII complex transits through a number of intracellular compartments in the endocytic pathway until it reaches the cell membrane for antigen presentation (PubMed:18305173, PubMed:9075930). Component of immunological synapses at the interface between T cell and APC (PubMed:15322540, PubMed:29884618).

### Tissue Location


Expressed in professional APCs: macrophages, dendritic cells and B cells (at protein level) (PubMed:15322540, PubMed:23783831, PubMed:31495665). Expressed in thymic epithelial cells (at protein level) (PubMed:23783831).

### Anti-HLA-DRA/Hla Dr Rabbit Monoclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


### Anti-HLA-DRA/Hla Dr Rabbit Monoclonal Antibody - Images




All lanes use the Antibody at 1:5K dilution for 1 hour at room temperature.




All lanes use the Antibody at 1:5K dilution for 1 hour at room temperature.




All lanes use the Antibody at 1:5K dilution for 1 hour at room temperature.



Western blot analysis of HLA-DRA expression in Raji cell lysate.



Immunohistochemical analysis of paraffin-embedded human lung, using HLA-DRA Antibody.



Immunofluorescent analysis of Raji cells, using HLA-DRA Antibody.