

**Anti-Aurora A AURKA Rabbit Monoclonal Antibody**  
Catalog # ABO13922**Specification****Anti-Aurora A AURKA Rabbit Monoclonal Antibody - Product Information**

Application	WB, IF, ICC, FC
Primary Accession	<a href="#">O14965</a>
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Human, Mouse
Clonality	Monoclonal
Format	Liquid

**Description**

Anti-Aurora A AURKA Rabbit Monoclonal Antibody . Tested in WB, ICC/IF, Flow Cytometry applications. This antibody reacts with Human, Mouse.

**Anti-Aurora A AURKA Rabbit Monoclonal Antibody - Additional Information**

**Gene ID** 6790

**Other Names**

Aurora kinase A {ECO:0000312|HGNC:HGNC:11393}, 2.7.11.1, Aurora 2, Aurora/IPL1-related kinase 1, ARK-1, Aurora-related kinase 1, Breast tumor-amplified kinase, Ipl1- and aurora-related kinase 1, AURKA ([HGNC:11393](http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=11393))

**Calculated MW**

45809 MW KDa

**Application Details**

WB 1:500-1:2000<br>ICC/IF 1:50-1:200<br>FC 1:<br>200

**Subcellular Localization**

Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, cytoskeleton, spindle pole. Cytoplasm, cytoskeleton, cilium basal body. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome, centriole. Detected at the neurite hillock in developing neurons (By similarity). Localizes at the centrosome in mitotic cells from early prophase until telophase, but also localizes to the spindle pole MTs from prophase to anaphase (PubMed:9606188, PubMed:17229885, PubMed:21225229). Colocalized with SIRT2 at centrosome (PubMed:22014574). Moves to the midbody during both telophase and cytokinesis (PubMed:17726514). Associates with both the pericentriolar material (PCM) and centrioles (PubMed:22014574)..

**Tissue Specificity**

Highly expressed in testis and weakly in skeletal muscle, thymus and spleen. Also highly expressed in colon, ovarian, prostate, neuroblastoma, breast and cervical cancer cell lines.

**Contents**

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

### Immunogen

A synthesized peptide derived from human Aurora A

### Purification

Affinity-chromatography

### Storage

**Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.**

## Anti-Aurora A AURKA Rabbit Monoclonal Antibody - Protein Information

Name AURKA ([HGNC:11393](#))

### Function

Mitotic serine/threonine kinase that contributes to the regulation of cell cycle progression (PubMed: [11039908](http://www.uniprot.org/citations/11039908)), PubMed: [12390251](http://www.uniprot.org/citations/12390251), PubMed: [17125279](http://www.uniprot.org/citations/17125279), PubMed: [17360485](http://www.uniprot.org/citations/17360485), PubMed: [18615013](http://www.uniprot.org/citations/18615013), PubMed: [26246606](http://www.uniprot.org/citations/26246606)). Associates with the centrosome and the spindle microtubules during mitosis and plays a critical role in various mitotic events including the establishment of mitotic spindle, centrosome duplication, centrosome separation as well as maturation, chromosomal alignment, spindle assembly checkpoint, and cytokinesis (PubMed: [14523000](http://www.uniprot.org/citations/14523000), PubMed: [26246606](http://www.uniprot.org/citations/26246606)). Required for normal spindle positioning during mitosis and for the localization of NUMA1 and DCTN1 to the cell cortex during metaphase (PubMed: [27335426](http://www.uniprot.org/citations/27335426)). Required for initial activation of CDK1 at centrosomes (PubMed: [13678582](http://www.uniprot.org/citations/13678582), PubMed: [15128871](http://www.uniprot.org/citations/15128871)). Phosphorylates numerous target proteins, including ARHGEF2, BORA, BRCA1, CDC25B, DLGP5, HDAC6, KIF2A, LATS2, NDEL1, PARD3, PPP1R2, PLK1, RASSF1, TACC3, p53/TP53 and TPX2 (PubMed: [11551964](http://www.uniprot.org/citations/11551964), PubMed: [14702041](http://www.uniprot.org/citations/14702041), PubMed: [15128871](http://www.uniprot.org/citations/15128871), PubMed: [15147269](http://www.uniprot.org/citations/15147269), PubMed: [15987997](http://www.uniprot.org/citations/15987997), PubMed: [17604723](http://www.uniprot.org/citations/17604723), PubMed: [18056443](http://www.uniprot.org/citations/18056443), PubMed: [18615013](http://www.uniprot.org/citations/18615013)). Regulates KIF2A tubulin depolymerase activity (PubMed: [19351716](http://www.uniprot.org/citations/19351716)). Important for microtubule formation and/or stabilization (PubMed: [18056443](http://www.uniprot.org/citations/18056443)). Required for normal axon formation (PubMed: [19812038](http://www.uniprot.org/citations/19812038)). Plays a role in microtubule remodeling during neurite extension (PubMed: [19668197](http://www.uniprot.org/citations/19668197)). Also acts as a key regulatory component of the p53/TP53 pathway, and particularly the checkpoint- response pathways critical for oncogenic transformation of cells, by phosphorylating

and destabilizing p53/TP53 (PubMed:<a href="http://www.uniprot.org/citations/14702041" target="\_blank">14702041</a>). Phosphorylates its own inhibitors, the protein phosphatase type 1 (PP1) isoforms, to inhibit their activity (PubMed:<a href="http://www.uniprot.org/citations/11551964" target="\_blank">11551964</a>). Inhibits cilia outgrowth (By similarity). Required for cilia disassembly via phosphorylation of HDAC6 and subsequent deacetylation of alpha-tubulin (PubMed:<a href="http://www.uniprot.org/citations/17604723" target="\_blank">17604723</a>, PubMed:<a href="http://www.uniprot.org/citations/20643351" target="\_blank">20643351</a>). Regulates protein levels of the anti-apoptosis protein BIRC5 by suppressing the expression of the SCF(FBXL7) E3 ubiquitin-protein ligase substrate adapter FBXL7 through the phosphorylation of the transcription factor FOXP1 (PubMed:<a href="http://www.uniprot.org/citations/28218735" target="\_blank">28218735</a>).

### Cellular Location

Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, cytoskeleton, spindle pole. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome, centriole {ECO:0000250|UniProtKB:P97477}. Cell projection, neuron projection {ECO:0000250|UniProtKB:P97477}. Cell projection, cilium. Cytoplasm, cytoskeleton, cilium basal body. Basolateral cell membrane {ECO:0000250|UniProtKB:F1PNY0}. Note=Detected at the neurite hillock in developing neurons (By similarity). Localizes at the centrosome in mitotic cells from early prophase until telophase, but also localizes to the spindle pole MTs from prophase to anaphase (PubMed:17229885, PubMed:21225229, PubMed:9606188). Colocalized with SIRT2 at centrosome (PubMed:22014574). Moves to the midbody during both telophase and cytokinesis (PubMed:17726514). Associates with both the pericentriolar material (PCM) and centrioles (PubMed:22014574). The localization to the spindle poles is regulated by AAAS (PubMed:26246606) {ECO:0000250|UniProtKB:P97477, ECO:0000269|PubMed:17229885, ECO:0000269|PubMed:17726514, ECO:0000269|PubMed:21225229, ECO:0000269|PubMed:22014574, ECO:0000269|PubMed:26246606, ECO:0000269|PubMed:9606188}

### Tissue Location


Highly expressed in testis and weakly in skeletal muscle, thymus and spleen. Also highly expressed in colon, ovarian, prostate, neuroblastoma, breast and cervical cancer cell lines


## Anti-Aurora A AURKA Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## Anti-Aurora A AURKA Rabbit Monoclonal Antibody - Images





Western blot analysis of Aurora A expression in HepG2 cell lysate.