

**Anti-Cyclin A2 Rabbit Monoclonal Antibody**  
Catalog # ABO13991**Specification****Anti-Cyclin A2 Rabbit Monoclonal Antibody - Product Information**

Application	WB, IHC, IF, ICC
Primary Accession	<a href="#">P20248</a>
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

**Description**

Anti-Cyclin A2 Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF applications. This antibody reacts with Human, Mouse, Rat.

**Anti-Cyclin A2 Rabbit Monoclonal Antibody - Additional Information**

**Gene ID** 890

**Other Names**

Cyclin-A2 {ECO:0000312|HGNC:HGNC:1578}, Cyclin-A, Cyclin A, CCNA2 ([http://www.genenames.org/cgi-bin/gene\\_symbol\\_report?hgnc\\_id=1578](http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=1578))

**Calculated MW**

48551 MW KDa

**Application Details**

WB 1:1000-1:2000<br>IHC 1:50-1:200<br>ICC/IF 1:50-1:200

**Subcellular Localization**

Nucleus. Cytoplasm. Cytoplasmic when associated with SCAPER.

**Contents**

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

**Immunogen**

A synthesized peptide derived from human Cyclin A2

**Purification**

Affinity-chromatography

**Storage**

**Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.**

## Anti-Cyclin A2 Rabbit Monoclonal Antibody - Protein Information

**Name** CCNA2 ([HGNC:1578](#))

### Function

Cyclin which controls both the G1/S and the G2/M transition phases of the cell cycle. Functions through the formation of specific serine/threonine protein kinase holoenzyme complexes with the cyclin- dependent protein kinases CDK1 or CDK2. The cyclin subunit confers the substrate specificity of these complexes and differentially interacts with and activates CDK1 and CDK2 throughout the cell cycle.

### Cellular Location

Nucleus. Cytoplasm. Note=Exclusively nuclear during interphase (PubMed:1312467). Detected in the nucleus and the cytoplasm at prophase (PubMed:1312467). Cytoplasmic when associated with SCAPER (PubMed:17698606).

## Anti-Cyclin A2 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## Anti-Cyclin A2 Rabbit Monoclonal Antibody - Images




Figure 1. Western blot analysis of Cyclin A2 using anti-Cyclin A2 antibody (M00700-1). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human Jurkat whole cell lysates,  
Lane 2: human Hela whole cell lysates,  
Lane 3: human K562 whole cell lysates,  
Lane 4: human MCF-7 whole cell lysates,  
Lane 5: rat PC-12 whole cell lysates,  
Lane 6: rat NRK whole cell lysates,  
Lane 7: mouse RAW264.7 whole cell lysates,

Lane 8: mouse ANA-1 whole cell lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Cyclin A2 antigen affinity purified monoclonal antibody (Catalog # M00700-1) at 1:500 overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:1000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for Cyclin A2 at approximately 55 kDa. The expected band size for Cyclin A2 is at 49 kDa.




Figure 2. IHC analysis of Cyclin A2 using anti-Cyclin A2 antibody (M00700-1).

Cyclin A2 was detected in a paraffin-embedded section of human colorectal adenocarcinoma tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1:50 rabbit anti-Cyclin A2 Antibody (M00700-1) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.




Figure 3. IHC analysis of Cyclin A2 using anti-Cyclin A2 antibody (M00700-1).

Cyclin A2 was detected in a paraffin-embedded section of human liver cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1:50 rabbit anti-Cyclin A2 Antibody (M00700-1) overnight at 4°C. Peroxidase Conjugated Goat

Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.




Figure 4. IHC analysis of Cyclin A2 using anti-Cyclin A2 antibody (M00700-1). Cyclin A2 was detected in a paraffin-embedded section of human lung cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1:50 rabbit anti-Cyclin A2 Antibody (M00700-1) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.






Figure 5. IHC analysis of Cyclin A2 using anti-Cyclin A2 antibody (M00700-1). Cyclin A2 was detected in a paraffin-embedded section of human spleen tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1:50 rabbit anti-Cyclin A2 Antibody (M00700-1) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.

Hela



Immunofluorescent analysis using the Antibody at 1:50 dilution.

Hela



Immunofluorescent analysis using the Antibody at 1:50 dilution.