

**Anti-IKK gamma IKBKG Rabbit Monoclonal Antibody**  
Catalog # ABO14007**Specification****Anti-IKK gamma IKBKG Rabbit Monoclonal Antibody - Product Information**

Application	WB, IHC, IF, ICC, IP, FC
Primary Accession	<a href="#">Q9Y6K9</a>
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

**Description**

Anti-IKK gamma IKBKG Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

**Anti-IKK gamma IKBKG Rabbit Monoclonal Antibody - Additional Information**

**Gene ID** 8517

**Other Names**

NF-kappa-B essential modulator, NEMO, FIP-3, Ikb kinase-associated protein 1, IKKAP1, Inhibitor of nuclear factor kappa-B kinase subunit gamma, I-kappa-B kinase subunit gamma, IKK-gamma, IKKG, Ikb kinase subunit gamma, NF-kappa-B essential modifier, IKBKG ([HGNC:5961](http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=5961)), FIP3, NEMO

**Calculated MW**

48198 MW KDa

**Application Details**

WB 1:500-1:2000<br>IHC 1:50-1:200<br>ICC/IF 1:50-1:200<br>IP 1:50<br>FC 1:50

**Subcellular Localization**

Cytoplasm. Nucleus. Sumoylated NEMO accumulates in the nucleus in response to genotoxic stress.

**Tissue Specificity**

Heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas.

**Contents**

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

**Immunogen**

A synthesized peptide derived from human IKK gamma

**Purification**

Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

## Anti-IKK gamma IKBKG Rabbit Monoclonal Antibody - Protein Information

Name IKBKG ([HGNC:5961](#))

Synonyms FIP3, NEMO

### Function

Regulatory subunit of the IKK core complex which phosphorylates inhibitors of NF-kappa-B thus leading to the dissociation of the inhibitor/NF-kappa-B complex and ultimately the degradation of the inhibitor (PubMed: [14695475](http://www.uniprot.org/citations/14695475)), PubMed: [20724660](http://www.uniprot.org/citations/20724660), PubMed: [21518757](http://www.uniprot.org/citations/21518757), PubMed: [9751060](http://www.uniprot.org/citations/9751060)). Its binding to scaffolding polyubiquitin plays a key role in IKK activation by multiple signaling receptor pathways (PubMed: [16547522](http://www.uniprot.org/citations/16547522), PubMed: [18287044](http://www.uniprot.org/citations/18287044), PubMed: [19033441](http://www.uniprot.org/citations/19033441), PubMed: [19185524](http://www.uniprot.org/citations/19185524), PubMed: [21606507](http://www.uniprot.org/citations/21606507), PubMed: [27777308](http://www.uniprot.org/citations/27777308), PubMed: [33567255](http://www.uniprot.org/citations/33567255)). Can recognize and bind both 'Lys-63'-linked and linear polyubiquitin upon cell stimulation, with a much higher affinity for linear polyubiquitin (PubMed: [16547522](http://www.uniprot.org/citations/16547522), PubMed: [18287044](http://www.uniprot.org/citations/18287044), PubMed: [19033441](http://www.uniprot.org/citations/19033441), PubMed: [19185524](http://www.uniprot.org/citations/19185524), PubMed: [21606507](http://www.uniprot.org/citations/21606507), PubMed: [27777308](http://www.uniprot.org/citations/27777308)). Could be implicated in NF-kappa-B-mediated protection from cytokine toxicity. Essential for viral activation of IRF3 (PubMed: [19854139](http://www.uniprot.org/citations/19854139)). Involved in TLR3- and IFIH1-mediated antiviral innate response; this function requires 'Lys- 27'-linked polyubiquitination (PubMed: [20724660](http://www.uniprot.org/citations/20724660)).

### Cellular Location

Cytoplasm. Nucleus Note=Sumoylated NEMO accumulates in the nucleus in response to genotoxic stress.

### Tissue Location

Heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas


## Anti-IKK gamma IKBKG Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)

- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Anti-IKK gamma IKBKG Rabbit Monoclonal Antibody - Images



Western blot analysis of IKK gamma expression in Jurkat cell lysate.



Immunohistochemical analysis of paraffin-embedded mouse colon, using IKK gamma Antibody.