

Anti-Aconitase 1 ACO1 Rabbit Monoclonal Antibody
Catalog # ABO14061

Specification

Anti-Aconitase 1 ACO1 Rabbit Monoclonal Antibody - Product Information

Application	WB, IF, ICC
Primary Accession	P21399
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

Description

Anti-Aconitase 1 ACO1 Rabbit Monoclonal Antibody . Tested in WB, ICC/IF applications. This antibody reacts with Human, Mouse, Rat.

Anti-Aconitase 1 ACO1 Rabbit Monoclonal Antibody - Additional Information

Gene ID 48

Other Names

Cytoplasmic aconitate hydratase, Aconitase, 4.2.1.3, Citrate hydro-lyase, Ferritin repressor protein, Iron regulatory protein 1, IRP1, Iron-responsive element-binding protein 1, IRE-BP 1, ACO1, IREB1

Calculated MW

98399 MW KDa

Application Details

WB 1:500-1:2000
ICC/IF 1:50-1:200

Subcellular Localization

Cytoplasm.

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human Aconitase 1

Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Anti-Aconitase 1 ACO1 Rabbit Monoclonal Antibody - Protein Information

Name ACO1

Synonyms IREB1

Function

Bifunctional iron sensor that switches between 2 activities depending on iron availability (PubMed:1281544, PubMed:1946430, PubMed:8041788). Iron deprivation, promotes its mRNA binding activity through which it regulates the expression of genes involved in iron uptake, sequestration and utilization (PubMed:1281544, PubMed:1946430, PubMed:23891004, PubMed:8041788). Binds to iron-responsive elements (IRES) in the untranslated region of target mRNAs preventing for instance the translation of ferritin and aminolevulinic acid synthase and stabilizing the transferrin receptor mRNA (PubMed:1281544, PubMed:1946430, PubMed:23891004, PubMed:8041788).

Cellular Location


Cytoplasm, cytosol.


Anti-Aconitase 1 ACO1 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Aconitase 1 ACO1 Rabbit Monoclonal Antibody - Images






All lanes use the Antibody at 1:5K dilution for 1 hour at room temperature.



All lanes use the Antibody at 1:5K dilution for 1 hour at room temperature.



Western blot analysis of Aconitase 1 in (1) mouse kidney lysate; (2) HepG2 cell lysate.