

Anti-CD3 epsilon Rabbit Monoclonal Antibody Catalog # ABO14071

Specification

Anti-CD3 epsilon Rabbit Monoclonal Antibody - Product Information

Application	IHC, IF, ICC, IP
Primary Accession	P07766
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid

Description

Anti-CD3 epsilon Rabbit Monoclonal Antibody . Tested in IHC, ICC/IF, IP applications. This antibody reacts with Human.

Anti-CD3 epsilon Rabbit Monoclonal Antibody - Additional Information

Gene ID 916

Other Names

T-cell surface glycoprotein CD3 epsilon chain, T-cell surface antigen T3/Leu-4 epsilon chain, CD3e, CD3E, T3E

Calculated MW

23147 MW KDa

Application Details

IHC 1:50-1:200
ICC/IF 1:50-1:200
IP 1:50

Subcellular Localization

Membrane; Single-pass type I membrane protein.

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human CD3 epsilon

Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Anti-CD3 epsilon Rabbit Monoclonal Antibody - Protein Information

Name CD3E

Synonyms T3E

Function

Part of the TCR-CD3 complex present on T-lymphocyte cell surface that plays an essential role in adaptive immune response. When antigen presenting cells (APCs) activate T-cell receptor (TCR), TCR- mediated signals are transmitted across the cell membrane by the CD3 chains CD3D, CD3E, CD3G and CD3Z. All CD3 chains contain immunoreceptor tyrosine-based activation motifs (ITAMs) in their cytoplasmic domain. Upon TCR engagement, these motifs become phosphorylated by Src family protein tyrosine kinases LCK and FYN, resulting in the activation of downstream signaling pathways (PubMed:2470098). In addition of this role of signal transduction in T-cell activation, CD3E plays an essential role in correct T-cell development. Initiates the TCR-CD3 complex assembly by forming the two heterodimers CD3D/CD3E and CD3G/CD3E. Participates also in internalization and cell surface down- regulation of TCR-CD3 complexes via endocytosis sequences present in CD3E cytosolic region (PubMed:10384095, PubMed:26507128).

Cellular Location


Cell membrane; Single-pass type I membrane protein

Anti-CD3 epsilon Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-CD3 epsilon Rabbit Monoclonal Antibody - Images



Immunohistochemical analysis of paraffin-embedded mouse spleen, using CD3 epsilon Antibody.