

Anti-AMPK beta 1 PRKAB1 Rabbit Monoclonal Antibody

Catalog # ABO14089

Specification

Anti-AMPK beta 1 PRKAB1 Rabbit Monoclonal Antibody - Product Information

ApplicationWB, FCPrimary AccessionO9Y478HostRabbitIsotypeRabbit IgGReactivityRat, Human, MouseClonalityMonoclonalFormatLiquidDescriptionAnti-AMPK beta 1 PRKAB1 Rabbit Monoclonal Antibody . Tested in WB, FlapplicationsThis aptibady reasts with HumanAnti-AMPK beta 1 PRKAB1 Rabbit Monoclonal Antibody . Tested in WB, Fl

Anti-AMPK beta 1 PRKAB1 Rabbit Monoclonal Antibody . Tested in WB, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

Anti-AMPK beta 1 PRKAB1 Rabbit Monoclonal Antibody - Additional Information

Gene ID 5564

Other Names 5'-AMP-activated protein kinase subunit beta-1, AMPK subunit beta-1, AMPKb, PRKAB1, AMPK

Calculated MW 30382 MW KDa

Application Details WB 1:1000-1:2000
FC 1:50

Contents Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen A synthesized peptide derived from human AMPK beta 1

Purification Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Anti-AMPK beta 1 PRKAB1 Rabbit Monoclonal Antibody - Protein Information

Name PRKAB1



Synonyms AMPK

Function


Non-catalytic subunit of AMP-activated protein kinase (AMPK), an energy sensor protein kinase that plays a key role in regulating cellular energy metabolism. In response to reduction of intracellular ATP levels, AMPK activates energy-producing pathways and inhibits energy-consuming processes: inhibits protein, carbohydrate and lipid biosynthesis, as well as cell growth and proliferation. AMPK acts via direct phosphorylation of metabolic enzymes, and by longer-term effects via phosphorylation of transcription regulators. Also acts as a regulator of cellular polarity by remodeling the actin cytoskeleton; probably by indirectly activating myosin. Beta non-catalytic subunit acts as a scaffold on which the AMPK complex assembles, via its Cterminus that bridges alpha (PRKAA1 or PRKAA2) and gamma subunits (PRKAG1, PRKAG2 or PRKAG3).

Anti-AMPK beta 1 PRKAB1 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Anti-AMPK beta 1 PRKAB1 Rabbit Monoclonal Antibody - Images



Western blot analysis of AMPK beta 1 expression in HeLa cell lysate.