

## Anti-Hexokinase 1 HK1 Rabbit Monoclonal Antibody Catalog # ABO14121

### Specification

---

#### Anti-Hexokinase 1 HK1 Rabbit Monoclonal Antibody - Product Information

Application	WB, IHC, IF, ICC, FC
Primary Accession	<a href="#">P19367</a>
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

#### Description

Anti-Hexokinase 1 HK1 Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

#### Anti-Hexokinase 1 HK1 Rabbit Monoclonal Antibody - Additional Information

Gene ID 3098

#### Other Names

Hexokinase-1, 2.7.1.1, Brain form hexokinase, Hexokinase type I, HK I, Hexokinase-A, HK1 ([HGNC:4922](http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=4922))

#### Calculated MW

102486 MW KDa

#### Application Details

WB 1:500-1:2000<br>IHC 1:50-1:200<br>ICC/IF 1:50-1:200<br>FC 1:200

#### Subcellular Localization

Mitochondrion outer membrane. Its hydrophobic N-terminal sequence may be involved in membrane binding.

#### Tissue Specificity

Isoform 2 is erythrocyte specific. Isoform 3 and isoform 4 are testis-specific.

#### Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

#### Immunogen

A synthesized peptide derived from human Hexokinase 1

#### Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term

**storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.**

## **Anti-Hexokinase 1 HK1 Rabbit Monoclonal Antibody - Protein Information**

**Name** HK1 ([HGNC:4922](#))

### **Function**

Catalyzes the phosphorylation of various hexoses, such as D- glucose, D-glucosamine, D-fructose, D-mannose and 2-deoxy-D-glucose, to hexose 6-phosphate (D-glucose 6-phosphate, D-glucosamine 6-phosphate, D-fructose 6-phosphate, D-mannose 6-phosphate and 2-deoxy-D-glucose 6- phosphate, respectively) (PubMed:<a href="http://www.uniprot.org/citations/1637300" target="\_blank">1637300</a>, PubMed:<a href="http://www.uniprot.org/citations/25316723" target="\_blank">25316723</a>, PubMed:<a href="http://www.uniprot.org/citations/27374331" target="\_blank">27374331</a>). Does not phosphorylate N-acetyl-D-glucosamine (PubMed:<a href="http://www.uniprot.org/citations/27374331" target="\_blank">27374331</a>). Mediates the initial step of glycolysis by catalyzing phosphorylation of D-glucose to D-glucose 6-phosphate (By similarity). Involved in innate immunity and inflammation by acting as a pattern recognition receptor for bacterial peptidoglycan (PubMed:<a href="http://www.uniprot.org/citations/27374331" target="\_blank">27374331</a>). When released in the cytosol, N-acetyl-D-glucosamine component of bacterial peptidoglycan inhibits the hexokinase activity of HK1 and causes its dissociation from mitochondrial outer membrane, thereby activating the NLRP3 inflammasome (PubMed:<a href="http://www.uniprot.org/citations/27374331" target="\_blank">27374331</a>).

### **Cellular Location**

Mitochondrion outer membrane; Peripheral membrane protein. Cytoplasm, cytosol. Note=The mitochondrial-binding peptide (MBP) region promotes association with the mitochondrial outer membrane (Probable). Dissociates from the mitochondrial outer membrane following inhibition by N-acetyl-D-glucosamine, leading to relocation to the cytosol (PubMed:27374331).

### **Tissue Location**


Isoform 2: Erythrocyte specific (Ref.6). Isoform 3: Testis-specific (PubMed:10978502). Isoform 4: Testis-specific (PubMed:10978502). {ECO:0000269|PubMed:10978502, ECO:0000269|Ref.6}


## **Anti-Hexokinase 1 HK1 Rabbit Monoclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## **Anti-Hexokinase 1 HK1 Rabbit Monoclonal Antibody - Images**







All lanes use the Antibody at 1:1W dilution for 1 hour at room temperature.




All lanes use the Antibody at 1:1W dilution for 1 hour at room temperature.




All lanes use the Antibody at 1:1W dilution for 1 hour at room temperature.



All lanes use the Antibody at 1:1W dilution for 1 hour at room temperature.



Western blot analysis of Hexokinase 1 expression in (1) MCF-7 cell lysate; (2) 293T cell lysate.



Immunohistochemical analysis of paraffin-embedded human lung carcinoma, using Hexokinase 1 Antibody.