

**Anti-S100 alpha 6 S100A6 Rabbit Monoclonal Antibody**  
Catalog # ABO14131**Specification****Anti-S100 alpha 6 S100A6 Rabbit Monoclonal Antibody - Product Information**

|                   |                          |
|-------------------|--------------------------|
| Application       | WB, IHC, IF, ICC, IP, FC |
| Primary Accession | <a href="#">P06703</a>   |
| Host              | Rabbit                   |
| Isotype           | Rabbit IgG               |
| Reactivity        | Rat, Human, Mouse        |
| Clonality         | Monoclonal               |
| Format            | Liquid                   |

**Description**

Anti-S100 alpha 6 S100A6 Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

**Anti-S100 alpha 6 S100A6 Rabbit Monoclonal Antibody - Additional Information**

**Gene ID** 6277

**Other Names**

Protein S100-A6, Calcyclin, Growth factor-inducible protein 2A9, MLN 4, Prolactin receptor-associated protein, PRA, S100 calcium-binding protein A6, S100A6, CACY

**Calculated MW**

10180 MW KDa

**Application Details**

WB 1:500-1:2000<br>IHC 1:50-1:200<br>ICC/IF 1:50-1:200<br>IP 1:50<br>FC 1:50

**Subcellular Localization**

Nucleus envelope. Cytoplasm. Cell membrane; Peripheral membrane protein; Cytoplasmic side.

**Contents**

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

**Immunogen**

A synthesized peptide derived from human S100 alpha 6

**Purification**

Affinity-chromatography

Storage

**Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.**

**Anti-S100 alpha 6 S100A6 Rabbit Monoclonal Antibody - Protein Information**

**Name** S100A6

**Synonyms** CACY

**Function**

May function as calcium sensor and modulator, contributing to cellular calcium signaling. May function by interacting with other proteins, such as TPR-containing proteins, and indirectly play a role in many physiological processes such as the reorganization of the actin cytoskeleton and in cell motility. Binds 2 calcium ions. Calcium binding is cooperative.

**Cellular Location**

Nucleus envelope. Cytoplasm. Cell membrane; Peripheral membrane protein; Cytoplasmic side

**Anti-S100 alpha 6 S100A6 Rabbit Monoclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Anti-S100 alpha 6 S100A6 Rabbit Monoclonal Antibody - Images**




Figure 1. Western blot analysis of S100A6 using anti-S100A6 antibody (M02043).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human Hela whole cell lysates,

Lane 2: human SW620 whole cell lysates,

Lane 3: human placenta tissue lysates,

Lane 4: mouse NIH/3T3 whole cell lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90

minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-S100A6 antigen affinity purified monoclonal antibody (Catalog # M02043) at 1:500 overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:500 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for S100A6 at approximately 10 kDa. The expected band size for S100A6 is at 10 kDa.