

**Anti-Androgen Receptor AR Rabbit Monoclonal Antibody**  
Catalog # ABO14233

**Specification**

**Anti-Androgen Receptor AR Rabbit Monoclonal Antibody - Product Information**

|                   |                        |
|-------------------|------------------------|
| Application       | WB, IHC, IF, ICC, IP   |
| Primary Accession | <a href="#">P10275</a> |
| Host              | Rabbit                 |
| Isotype           | Rabbit IgG             |
| Reactivity        | Human                  |
| Clonality         | Monoclonal             |
| Format            | Liquid                 |

**Description**

Anti-Androgen Receptor AR Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, IP applications. This antibody reacts with Human.

**Anti-Androgen Receptor AR Rabbit Monoclonal Antibody - Additional Information**

**Gene ID** 367

**Other Names**

Androgen receptor, Dihydrotestosterone receptor, Nuclear receptor subfamily 3 group C member 4, AR, DHTR, NR3C4

**Calculated MW**

98989 MW KDa

**Application Details**

WB 1:500-1:2000<br>IHC 1:50-1:100<br>ICC/IF 1:50-1:100<br>IP 1:30

**Subcellular Localization**

Nucleus. Cytoplasm. Predominantly cytoplasmic in unligated form but translocates to the nucleus upon ligand-binding. Can also translocate to the nucleus in unligated form in the presence of GNB2L1.

**Tissue Specificity**

Isoform 2 is mainly expressed in heart and skeletal muscle..

**Contents**

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

**Immunogen**

A synthesized peptide derived from human Androgen Receptor

**Purification**

Affinity-chromatography

**Storage**

**Store at -20°C for one year. For short term**

storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

## Anti-Androgen Receptor AR Rabbit Monoclonal Antibody - Protein Information

**Name** AR

**Synonyms** DHTR, NR3C4

### Function

Steroid hormone receptors are ligand-activated transcription factors that regulate eukaryotic gene expression and affect cellular proliferation and differentiation in target tissues (PubMed:<a href="http://www.uniprot.org/citations/19022849" target="\_blank">19022849</a>). Transcription factor activity is modulated by bound coactivator and corepressor proteins like ZBTB7A that recruits NCOR1 and NCOR2 to the androgen response elements/ARE on target genes, negatively regulating androgen receptor signaling and androgen-induced cell proliferation (PubMed:<a href="http://www.uniprot.org/citations/20812024" target="\_blank">20812024</a>). Transcription activation is also down-regulated by NROB2. Activated, but not phosphorylated, by HIPK3 and ZIPK/DAPK3.

### Cellular Location

Nucleus. Cytoplasm Note=Detected at the promoter of target genes (PubMed:25091737) Predominantly cytoplasmic in unligated form but translocates to the nucleus upon ligand-binding. Can also translocate to the nucleus in unligated form in the presence of RACK1.

### Tissue Location


[Isoform 2]: Mainly expressed in heart and skeletal muscle.


## Anti-Androgen Receptor AR Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## Anti-Androgen Receptor AR Rabbit Monoclonal Antibody - Images







All lanes use the Antibody at 1:5K dilution for 1 hour at room temperature.




All lanes use the Antibody at 1:5K dilution for 1 hour at room temperature.




All lanes use the Antibody at 1:5K dilution for 1 hour at room temperature.




All lanes use the Antibody at 1:5K dilution for 1 hour at room temperature.




All lanes use the Antibody at 1:5K dilution for 1 hour at room temperature.




All lanes use the Antibody at 1:5K dilution for 1 hour at room temperature.




Western blot analysis of Androgen Receptor expression in (1) T47D cell lysate;(2) LnCaP cell lysate.




Immunohistochemical analysis of paraffin-embedded human prostate carcinoma, using Androgen Receptor Antibody.



Immunofluorescent analysis using the Antibody at 1:50 dilution.



Immunofluorescent analysis using the Antibody at 1:150 dilution.



Immunofluorescent analysis of Lncap cells, using Androgen Receptor Antibody.