

Anti-Apolipoprotein A I APOA1 Antibody Picoband™ (monoclonal, 17G5)
Catalog # ABO14241

Specification

Anti-Apolipoprotein A I APOA1 Antibody Picoband™ (monoclonal, 17G5) - Product Information

Application	E
Primary Accession	P02647
Host	Mouse
Isotype	Mouse IgG1
Reactivity	Human
Clonality	Monoclonal
Format	Lyophilized

Description

Anti-Apolipoprotein A I APOA1 Antibody Picoband™ (monoclonal, 17G5) . Tested in ELISA applications. This antibody reacts with Human.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500 µg/ml.

Anti-Apolipoprotein A I APOA1 Antibody Picoband™ (monoclonal, 17G5) - Additional Information

Gene ID 335

Other Names

Apolipoprotein A-I, Apo-AI, ApoA-I, Apolipoprotein A1, Proapolipoprotein A-I, ProapoA-I, Truncated apolipoprotein A-I, Apolipoprotein A-I(1-242), APOA1 ([HGNC:600](http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=600))

Application Details

ELISA (Cap), 1-5 µg/ml

Subcellular Localization

Secreted.

Tissue Specificity

Major protein of plasma HDL, also found in chylomicrons. Synthesized in the liver and small intestine. The oxidized form at Met-110 and Met-136 is increased in individuals with increased risk for coronary artery disease, such as in carrier of the eNOSa/b genotype and exposure to cigarette smoking. It is also present in increased levels in aortic lesions relative to native ApoA-I and increased levels are seen with increasing severity of disease.

Contents

Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.

Immunogen

E. coli-derived mouse Apolipoprotein A I recombinant protein (Position: D25-Q264). Mouse

Apolipoprotein A I shares 64% and 68.6% amino acid (aa) sequence identity with human and rat Apolipoprotein A I, respectively.

Purification

Affinity-chromatography

Cross Reactivity

No cross-reactivity with other proteins.

Storage

Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.

Anti-Apolipoprotein A I APOA1 Antibody Picoband™ (monoclonal, 17G5) - Protein Information

Name APOA1 ([HGNC:600](#))

Function

Participates in the reverse transport of cholesterol from tissues to the liver for excretion by promoting cholesterol efflux from tissues and by acting as a cofactor for the lecithin cholesterol acyltransferase (LCAT). As part of the SPAP complex, activates spermatozoa motility.

Cellular Location

Secreted.

Tissue Location

Major protein of plasma HDL, also found in chylomicrons. Synthesized in the liver and small intestine. The oxidized form at Met-110 and Met-136 is increased in individuals with increased risk for coronary artery disease, such as in carrier of the eNOSa/b genotype and exposure to cigarette smoking. It is also present in increased levels in aortic lesions relative to native ApoA-I and increased levels are seen with increasing severity of disease

Anti-Apolipoprotein A I APOA1 Antibody Picoband™ (monoclonal, 17G5) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Apolipoprotein A I APOA1 Antibody Picoband™ (monoclonal, 17G5) - Images

Anti-Apolipoprotein A I APOA1 Antibody Picoband™ (monoclonal, 17G5) - Background

Apolipoprotein A-1, also known as APOA1, is a human protein with a specific role in lipid metabolism. It binds to lipopolysaccharide or endotoxin, and has a major role in the anti-endotoxin function of HDL. The gene is mapped to 11q23. And it is a single polypeptide chain with 243 amino

acid residues of known primary amino acid sequence. The ApoA-I protein promotes cholesterol efflux from tissues to the liver for excretion. It is a cofactor for lecithin cholesterolacyltransferase (LCAT) which is responsible for the formation of most plasma cholesteryl esters. ApoA-I is also isolated as a prostacyclin (PGI₂) stabilizing factor, and thus may have an anticlotting effect. Defects in the gene encoding it are associated with HDL deficiencies, including Tangier disease, and with systemic non-neuropathic amyloidosis. Additionally, ApoA-I overexpression promotes macrophage-specific reverse cholesterol transport.