

**Anti-Vitamin D Receptor VDR Rabbit Monoclonal Antibody**  
Catalog # ABO14260**Specification****Anti-Vitamin D Receptor VDR Rabbit Monoclonal Antibody - Product Information**

Application	WB, IP
Primary Accession	<a href="#">P11473</a>
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

**Description**

Anti-Vitamin D Receptor VDR Rabbit Monoclonal Antibody . Tested in WB, IP applications. This antibody reacts with Human, Mouse, Rat.

**Anti-Vitamin D Receptor VDR Rabbit Monoclonal Antibody - Additional Information**

**Gene ID** 7421

**Other Names**

Vitamin D3 receptor, VDR, 1, 25-dihydroxyvitamin D3 receptor, Nuclear receptor subfamily 1 group I member 1, VDR ([http://www.genenames.org/cgi-bin/gene\\_symbol\\_report?hgnc\\_id=12679](http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=12679)), NR111

**Calculated MW**

48289 MW KDa

**Application Details**

WB 1:500-1:2000<br>IP 1:50

**Subcellular Localization**

Nucleus.

**Contents**

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

**Immunogen**

A synthesized peptide derived from human Vitamin D Receptor

**Purification**

Affinity-chromatography

**Storage**

**Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.**

## Anti-Vitamin D Receptor VDR Rabbit Monoclonal Antibody - Protein Information

Name VDR ([HGNC:12679](#))

Synonyms NR111

### Function

Nuclear receptor for calcitriol, the active form of vitamin D3 which mediates the action of this vitamin on cells (PubMed:[10678179](http://www.uniprot.org/citations/10678179), PubMed:[15728261](http://www.uniprot.org/citations/15728261), PubMed:[16913708](http://www.uniprot.org/citations/16913708), PubMed:[28698609](http://www.uniprot.org/citations/28698609)). Enters the nucleus upon vitamin D3 binding where it forms heterodimers with the retinoid X receptor/RXR (PubMed:[28698609](http://www.uniprot.org/citations/28698609)). The VDR-RXR heterodimers bind to specific response elements on DNA and activate the transcription of vitamin D3-responsive target genes (PubMed:[28698609](http://www.uniprot.org/citations/28698609)). Plays a central role in calcium homeostasis (By similarity). Also functions as a receptor for the secondary bile acid lithocholic acid (LCA) and its metabolites (PubMed:[12016314](http://www.uniprot.org/citations/12016314), PubMed:[32354638](http://www.uniprot.org/citations/32354638)).

### Cellular Location


Nucleus {ECO:0000255|PROSITE-ProRule:PRU00407, ECO:0000269|PubMed:12145331, ECO:0000269|PubMed:16207705, ECO:0000269|PubMed:28698609}. Cytoplasm Note=Localizes mainly to the nucleus (PubMed:12145331, PubMed:28698609). Translocated into the nucleus via both ligand- dependent and ligand-independent pathways; ligand-independent nuclear translocation is mediated by IPO4 (PubMed:16207705)


## Anti-Vitamin D Receptor VDR Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## Anti-Vitamin D Receptor VDR Rabbit Monoclonal Antibody - Images





All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.




Figure 1. Western blot analysis of VDR using anti-VDR antibody (M00210).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human HeLa whole cell lysates,  
 Lane 2: mouse kidney tissue lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-VDR antigen affinity purified monoclonal antibody (Catalog # M00210) at 1:500 overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for VDR at approximately 48 kDa. The expected band size for VDR is at 48 kDa.