

**Anti-Wilms Tumor Protein WT1 Rabbit Monoclonal Antibody**  
Catalog # ABO14264

**Specification**

---

**Anti-Wilms Tumor Protein WT1 Rabbit Monoclonal Antibody - Product Information**

Application	WB, IHC, IF, ICC, FC
Primary Accession	<a href="#">P19544</a>
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Human, Mouse
Clonality	Monoclonal
Format	Liquid

**Description**

Anti-Wilms Tumor Protein WT1 Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, Flow Cytometry applications. This antibody reacts with Human, Mouse.

**Anti-Wilms Tumor Protein WT1 Rabbit Monoclonal Antibody - Additional Information**

**Gene ID** 7490

**Other Names**

Wilms tumor protein, WT33, WT1

**Calculated MW**

49188 MW KDa

**Application Details**

WB 1:500-1:1000<br>IHC 1:50-1:200<br>ICC/IF 1:50-1:200<br>FC 1:50

**Subcellular Localization**

Nucleus. Nucleus, nucleolus. Cytoplasm. Shuttles between nucleus and cytoplasm..

**Tissue Specificity**

Expressed in the kidney and a subset of hematopoietic cells.

**Contents**

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

**Immunogen**

A synthesized peptide derived from human Wilms Tumor Protein

**Purification**

Affinity-chromatography

**Storage**

**Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.**

## Anti-Wilms Tumor Protein WT1 Rabbit Monoclonal Antibody - Protein Information

**Name** WT1

### Function

Transcription factor that plays an important role in cellular development and cell survival (PubMed:<a href="http://www.uniprot.org/citations/7862533" target="\_blank">7862533</a>). Recognizes and binds to the DNA sequence 5'-GCG(T/G)GGGCG-3' (PubMed:<a href="http://www.uniprot.org/citations/17716689" target="\_blank">17716689</a>, PubMed:<a href="http://www.uniprot.org/citations/25258363" target="\_blank">25258363</a>, PubMed:<a href="http://www.uniprot.org/citations/7862533" target="\_blank">7862533</a>). Regulates the expression of numerous target genes, including EPO. Plays an essential role for development of the urogenital system. It has a tumor suppressor as well as an oncogenic role in tumor formation. Function may be isoform-specific: isoforms lacking the KTS motif may act as transcription factors (PubMed:<a href="http://www.uniprot.org/citations/15520190" target="\_blank">15520190</a>). Isoforms containing the KTS motif may bind mRNA and play a role in mRNA metabolism or splicing (PubMed:<a href="http://www.uniprot.org/citations/16934801" target="\_blank">16934801</a>). Isoform 1 has lower affinity for DNA, and can bind RNA (PubMed:<a href="http://www.uniprot.org/citations/19123921" target="\_blank">19123921</a>).

### Cellular Location

Nucleus. Nucleus, nucleolus. Cytoplasm. Note=Isoforms lacking the KTS motif have a diffuse nuclear location (PubMed:15520190). Shuttles between nucleus and cytoplasm. {ECO:0000250, ECO:0000269|PubMed:15520190} [Isoform 4]: Nucleus, nucleoplasm

### Tissue Location


Expressed in the kidney and a subset of hematopoietic cells


## Anti-Wilms Tumor Protein WT1 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## Anti-Wilms Tumor Protein WT1 Rabbit Monoclonal Antibody - Images







All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.



All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.



Western blot analysis of WT1 expression in K562 cell lysate.



Immunohistochemical analysis of paraffin-embedded mouse kidney, using Wilms Tumor Protein Antibody.