

**Anti-Histone H4 (mono methyl K16) HIST1H4A Rabbit Monoclonal Antibody**  
Catalog # ABO14318

**Specification**

**Anti-Histone H4 (mono methyl K16) HIST1H4A Rabbit Monoclonal Antibody - Product Information**

|                   |                        |
|-------------------|------------------------|
| Application       | WB, IF, ICC            |
| Primary Accession | <a href="#">P62805</a> |
| Host              | Rabbit                 |
| Isotype           | Rabbit IgG             |
| Reactivity        | Human, Mouse           |
| Clonality         | Monoclonal             |
| Format            | Liquid                 |

**Description**

Anti-Histone H4 (mono methyl K16) HIST1H4A Rabbit Monoclonal Antibody . Tested in WB, ICC/IF applications. This antibody reacts with Human, Mouse.

**Anti-Histone H4 (mono methyl K16) HIST1H4A Rabbit Monoclonal Antibody - Additional Information**

**Gene ID** 121504;554313;8294;8359;8360;8361;8362;8363;8364;8365;8366;8367;8368;8370

**Other Names**

Histone H4, H4C1, H4/A, H4FA, HIST1H4A

**Calculated MW**

11367 MW KDa

**Application Details**

WB 1:500-1:2000<br>ICC/IF 1:500-1:2000

**Subcellular Localization**

Nucleus. Chromosome.

**Contents**

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

**Immunogen**

A synthesized peptide derived from human Histone H4 (mono methyl K16)

**Purification**

Affinity-chromatography

**Storage**

**Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.**

## Anti-Histone H4 (mono methyl K16) HIST1H4A Rabbit Monoclonal Antibody - Protein Information

**Name** H4C1

**Synonyms** H4/A, H4FA, HIST1H4A

### Function

Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.

### Cellular Location


Nucleus. Chromosome.

## Anti-Histone H4 (mono methyl K16) HIST1H4A Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## Anti-Histone H4 (mono methyl K16) HIST1H4A Rabbit Monoclonal Antibody - Images



Western blot analysis of Histone H4 (mono methyl K16) expression in (1) NIH/3T3 cell lysate; (2) A549 cell lysate.