

Anti-Collagen IV COL4A1 Antibody Picoband™ (monoclonal, 3G3)
Catalog # ABO14330**Specification****Anti-Collagen IV COL4A1 Antibody Picoband™ (monoclonal, 3G3) - Product Information**

Application	WB, IHC, IF
Primary Accession	P02462
Host	Mouse
Isotype	Mouse IgG1
Reactivity	Human
Clonality	Monoclonal
Format	Lyophilized

Description

Anti-Collagen IV COL4A1 Antibody Picoband™ (monoclonal, 3G3) . Tested in IF, IHC, WB applications. This antibody reacts with Human.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-Collagen IV COL4A1 Antibody Picoband™ (monoclonal, 3G3) - Additional Information**Gene ID 1282****Other Names**

Collagen alpha-1(IV) chain, Arresten, COL4A1 (http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=2202)
HGNC:2202

Calculated MW

220 kDa KDa

Application Details

Western blot, 0.1-0.5 µg/ml
 Immunohistochemistry (Paraffin-embedded Section), 0.5-1 µg/ml
 Immunofluorescence, 2 µg/ml

Subcellular Localization

Secreted, extracellular space, extracellular matrix, basement membrane

Tissue Specificity

Highly expressed in placenta.

Contents

Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.

Immunogen

E.coli-derived human Collagen IV recombinant protein (Position: G1445-T1669). Human Collagen IV shares 97% amino acid (aa) sequence identity with mouse Collagen IV.

Cross Reactivity

No cross-reactivity with other proteins.

Storage

Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.

Anti-Collagen IV COL4A1 Antibody Picoband™ (monoclonal, 3G3) - Protein Information

Name COL4A1 ([HGNC:2202](#))

Function

Type IV collagen is the major structural component of glomerular basement membranes (GBM), forming a 'chicken-wire' meshwork together with laminins, proteoglycans and entactin/nidogen.

Cellular Location

Secreted, extracellular space, extracellular matrix, basement membrane
{ECO:0000250|UniProtKB:P02463}

Tissue Location

Highly expressed in placenta.

Anti-Collagen IV COL4A1 Antibody Picoband™ (monoclonal, 3G3) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Collagen IV COL4A1 Antibody Picoband™ (monoclonal, 3G3) - Images




Figure 1. IHC analysis of Collagen IV using anti-Collagen IV antibody (M01411). Collagen IV was detected in paraffin-embedded section of human lung cancer tissue. Heat

mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 µg/ml mouse anti-Collagen IV Antibody (M01411) overnight at 4°C. Biotinylated goat anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC)(Catalog # SA1021) with DAB as the chromogen.




Figure 2. IHC analysis of Collagen IV using anti-Collagen IV antibody (M01411).

Collagen IV was detected in paraffin-embedded section of human placenta tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 µg/ml mouse anti-Collagen IV Antibody (M01411) overnight at 4°C. Biotinylated goat anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC)(Catalog # SA1021) with DAB as the chromogen.




Figure 3. Western blot analysis of Collagen IV using anti-Collagen IV antibody (M01411).

Electrophoresis was performed on a 8% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions.

Lane 1: human HEK293T whole cell lysate,

Lane 2: human Hela whole cell lysate,

Lane 3: human A549 whole cell lysate.

After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with mouse anti-Collagen IV antigen affinity purified monoclonal antibody (Catalog #

M01411) at 0.5 µg/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-mouse IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1001) with Tanon 5200 system. A specific band was detected for Collagen IV at approximately 220KD. The expected band size for Collagen IV is at 161KD.

Anti-Collagen IV COL4A1 Antibody Picoband™ (monoclonal, 3G3) - Background

COL4A1, also known as ICH or Collagen alpha-1 (IV), is a protein that in humans is encoded by the COL4A1 gene. It is mapped to 13q34. This gene encodes the major type IV alpha collagen chain of basement membranes. Like the other members of the type IV collagen gene family, this gene is organized in a head-to-head conformation with another type IV collagen gene so that each gene pair shares a common promoter. COL4A1 binds to alpha-1/beta-1 integrin and inhibits migration, proliferation, and tube formation by endothelial cells. It is also a potential therapeutic candidate for targeting tumor angiogenesis.