

## Anti-c-Myc Monoclonal Antibody Catalog # ABO14355

### Specification

---

#### Anti-c-Myc Monoclonal Antibody - Product Information

|                   |                          |
|-------------------|--------------------------|
| Application       | WB, IHC, IF, ICC, IP, FC |
| Primary Accession | <a href="#">P01106</a>   |
| Host              | Rabbit                   |
| Isotype           | Rabbit IgG               |
| Reactivity        | Rat, Human, Mouse        |
| Clonality         | Monoclonal               |
| Format            | Liquid                   |

#### Description

Anti-c-Myc Monoclonal Antibody . Tested in WB, IHC, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

#### Anti-c-Myc Monoclonal Antibody - Additional Information

Gene ID 4609

#### Other Names

Myc proto-oncogene protein, Class E basic helix-loop-helix protein 39, bHLHe39, Proto-oncogene c-Myc, Transcription factor p64, MYC, BHLHE39

#### Application Details

WB 1:500-1:2000<br>IHC 1:50-1:200<br>ICC/IF 1:50-1:200<br>IP 1:50<br>FC 1:50

#### Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

#### Immunogen

A synthesized peptide derived from human c-Myc Transcription factor that binds DNA in a non-specific manner, yet also specifically recognizes the core sequence 5'-CAC[GA]TG-3'. Activates the transcription of growth-related genes. Myc a proto-oncogenic transcription factor that plays a role in cell proliferation, apoptosis and in the development of human tumors.. Seems to activate the transcription of growth-related genes.

#### Purification

Affinity-chromatography

Storage

**Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.**

#### Anti-c-Myc Monoclonal Antibody - Protein Information

## Name MYC

## Synonyms BHLHE39

### Function

Transcription factor that binds DNA in a non-specific manner, yet also specifically recognizes the core sequence 5'-CAC[GA]TG-3' (PubMed:<a href="http://www.uniprot.org/citations/24940000" target="\_blank">24940000</a>, PubMed:<a href="http://www.uniprot.org/citations/25956029" target="\_blank">25956029</a>). Activates the transcription of growth-related genes (PubMed:<a href="http://www.uniprot.org/citations/24940000" target="\_blank">24940000</a>, PubMed:<a href="http://www.uniprot.org/citations/25956029" target="\_blank">25956029</a>). Binds to the VEGFA promoter, promoting VEGFA production and subsequent sprouting angiogenesis (PubMed:<a href="http://www.uniprot.org/citations/24940000" target="\_blank">24940000</a>, PubMed:<a href="http://www.uniprot.org/citations/25956029" target="\_blank">25956029</a>). Regulator of somatic reprogramming, controls self-renewal of embryonic stem cells (By similarity). Functions with TAF6L to activate target gene expression through RNA polymerase II pause release (By similarity). Positively regulates transcription of HNRNPA1, HNRNPA2 and PTBP1 which in turn regulate splicing of pyruvate kinase PKM by binding repressively to sequences flanking PKM exon 9, inhibiting exon 9 inclusion and resulting in exon 10 inclusion and production of the PKM M2 isoform (PubMed:<a href="http://www.uniprot.org/citations/20010808" target="\_blank">20010808</a>).

### Cellular Location


Nucleus, nucleoplasm. Nucleus, nucleolus. Nucleus. Cytoplasm


## Anti-c-Myc Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## Anti-c-Myc Monoclonal Antibody - Images







All lanes use the Antibody at 1:1W dilution for 1 hour at room temperature.




Immunofluorescent analysis using the Antibody at 1:50 dilution.



Western blot analysis of c-Myc expression in (1) Jurkat cell lysate;(2) K562 cell lysate.



Immunohistochemical analysis of paraffin-embedded human lung cancer, using c-Myc Antibody.



Immunofluorescent analysis of HeLa cells, using c-Myc Antibody.