

**Anti-SPHK1 Monoclonal Antibody**  
Catalog # ABO14420**Specification****Anti-SPHK1 Monoclonal Antibody - Product Information**

Application	WB, FC
Primary Accession	<a href="#">O9NYA1</a>
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

**Description**

Anti-SPHK1 Monoclonal Antibody . Tested in WB, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

**Anti-SPHK1 Monoclonal Antibody - Additional Information**

**Gene ID** 8877

**Other Names**

Sphingosine kinase 1, SK 1, SPK 1, 2.7.1.91, Acetyltransferase SPHK1, 2.3.1.-, SPHK1 ([http://www.genenames.org/cgi-bin/gene\\_symbol\\_report?hgnc\\_id=11240](http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=11240))  
HGNC:11240

**Application Details**

WB 1:500-1:2000  
FC 1:50

**Contents**

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

**Immunogen**

A synthesized peptide derived from human SPHK1

**Purification**

Affinity-chromatography

**Storage**

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

**Anti-SPHK1 Monoclonal Antibody - Protein Information**

**Name** SPHK1 ([HGNC:11240](#))

**Function**

Catalyzes the phosphorylation of sphingosine to form sphingosine 1-phosphate (SPP), a lipid mediator with both intra- and extracellular functions. Also acts on D-erythro-sphingosine and to a lesser extent sphinganine, but not other lipids, such as D,L-threo- dihydrosphingosine, N,N-dimethylsphingosine, diacylglycerol, ceramide, or phosphatidylinositol (PubMed:<a href="http://www.uniprot.org/citations/11923095" target="\_blank">11923095</a>, PubMed:<a href="http://www.uniprot.org/citations/20577214" target="\_blank">20577214</a>, PubMed:<a href="http://www.uniprot.org/citations/23602659" target="\_blank">23602659</a>, PubMed:<a href="http://www.uniprot.org/citations/24929359" target="\_blank">24929359</a>, PubMed:<a href="http://www.uniprot.org/citations/29662056" target="\_blank">29662056</a>). In contrast to proapoptotic SPHK2, has a negative effect on intracellular ceramide levels, enhances cell growth and inhibits apoptosis (PubMed:<a href="http://www.uniprot.org/citations/16118219" target="\_blank">16118219</a>). Involved in the regulation of inflammatory response and neuroinflammation. Via the product sphingosine 1-phosphate, stimulates TRAF2 E3 ubiquitin ligase activity, and promotes activation of NF- kappa-B in response to TNF signaling leading to IL17 secretion (PubMed:<a href="http://www.uniprot.org/citations/20577214" target="\_blank">20577214</a>). In response to TNF and in parallel to NF-kappa-B activation, negatively regulates RANTES induction through p38 MAPK signaling pathway (PubMed:<a href="http://www.uniprot.org/citations/23935096" target="\_blank">23935096</a>). Involved in endocytic membrane trafficking induced by sphingosine, recruited to dilate endosomes, also plays a role on later stages of endosomal maturation and membrane fusion independently of its kinase activity (PubMed:<a href="http://www.uniprot.org/citations/24929359" target="\_blank">24929359</a>, PubMed:<a href="http://www.uniprot.org/citations/28049734" target="\_blank">28049734</a>). In Purkinje cells, seems to be also involved in the regulation of autophagosome-lysosome fusion upon VEGFA (PubMed:<a href="http://www.uniprot.org/citations/25417698" target="\_blank">25417698</a>).

#### Cellular Location

Cytoplasm. Nucleus. Cell membrane. Endosome membrane; Peripheral membrane protein. Membrane, clathrin-coated pit. Synapse {ECO:0000250|UniProtKB:Q8CI15} Note=Translocated from the cytoplasm to the plasma membrane in a CIB1- dependent manner (PubMed:19854831). Binds to membranes containing negatively charged lipids but not neutral lipids (PubMed:24929359) Recruited to endocytic membranes by sphingosine where promotes membrane fusion (By similarity). {ECO:0000250|UniProtKB:Q8CI15, ECO:0000269|PubMed:19854831, ECO:0000269|PubMed:24929359}

#### Tissue Location


Widely expressed with highest levels in adult liver, kidney, heart and skeletal muscle. Expressed in brain cortex (at protein level) (PubMed:29662056).


#### Anti-SPHK1 Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### Anti-SPHK1 Monoclonal Antibody - Images






Western blot analysis of SPHK1 expression in Raji cell lysate.



All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.



All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.