

**Anti-HMGB2 Rabbit Monoclonal Antibody**  
Catalog # ABO14424**Specification**

---

**Anti-HMGB2 Rabbit Monoclonal Antibody - Product Information**

|                   |                        |
|-------------------|------------------------|
| Application       | WB, IHC, IF, ICC, IP   |
| Primary Accession | <a href="#">P26583</a> |
| Host              | Rabbit                 |
| Isotype           | Rabbit IgG             |
| Reactivity        | Rat, Human, Mouse      |
| Clonality         | Monoclonal             |
| Format            | Liquid                 |

**Description**

Anti-HMGB2 Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, IP applications. This antibody reacts with Human, Mouse, Rat.

**Anti-HMGB2 Rabbit Monoclonal Antibody - Additional Information**

**Gene ID** 3148

**Other Names**

High mobility group protein B2, High mobility group protein 2, HMG-2, HMGB2, HMG2

**Application Details**

WB 1:1000-1:5000<br>IHC 1:50-1:200<br>ICC/IF 1:50-1:200<br>IP 1:50

**Contents**

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

**Immunogen**

A synthesized peptide derived from human HMGB2

**Purification**

Affinity-chromatography

**Storage**

**Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.**

**Anti-HMGB2 Rabbit Monoclonal Antibody - Protein Information**

**Name** HMGB2

**Synonyms** HMG2

**Function**

Multifunctional protein with various roles in different cellular compartments. May act in a redox sensitive manner. In the nucleus is an abundant chromatin-associated non-histone protein involved in transcription, chromatin remodeling and V(D)J recombination and probably other processes. Binds DNA with a preference to non- canonical DNA structures such as single-stranded DNA. Can bent DNA and enhance DNA flexibility by looping thus providing a mechanism to promote activities on various gene promoters by enhancing transcription factor binding and/or bringing distant regulatory sequences into close proximity (PubMed:<a href="http://www.uniprot.org/citations/11909973" target="\_blank">11909973</a>, PubMed:<a href="http://www.uniprot.org/citations/18413230" target="\_blank">18413230</a>, PubMed:<a href="http://www.uniprot.org/citations/19522541" target="\_blank">19522541</a>, PubMed:<a href="http://www.uniprot.org/citations/19965638" target="\_blank">19965638</a>, PubMed:<a href="http://www.uniprot.org/citations/20123072" target="\_blank">20123072</a>, PubMed:<a href="http://www.uniprot.org/citations/7797075" target="\_blank">7797075</a>). Involved in V(D)J recombination by acting as a cofactor of the RAG complex: acts by stimulating cleavage and RAG protein binding at the 23 bp spacer of conserved recombination signal sequences (RSS) (By similarity). Proposed to be involved in the innate immune response to nucleic acids by acting as a promiscuous immunogenic DNA/RNA sensor which cooperates with subsequent discriminative sensing by specific pattern recognition receptors (By similarity). In the extracellular compartment acts as a chemokine. Promotes proliferation and migration of endothelial cells implicating AGER/RAGE (PubMed:<a href="http://www.uniprot.org/citations/19811285" target="\_blank">19811285</a>). Has antimicrobial activity in gastrointestinal epithelial tissues (PubMed:<a href="http://www.uniprot.org/citations/23877675" target="\_blank">23877675</a>). Involved in inflammatory response to antigenic stimulus coupled with pro- inflammatory activity (By similarity). Involved in modulation of neurogenesis probably by regulation of neural stem proliferation (By similarity). Involved in articular cartilage surface maintenance implicating LEF1 and the Wnt/beta-catenin pathway (By similarity).

#### Cellular Location

Nucleus. Chromosome. Cytoplasm. Secreted. Note=In basal state predominantly nuclear.

#### Tissue Location

Expressed in gastric and intestinal tissues (at protein level).


### Anti-HMGB2 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Anti-HMGB2 Rabbit Monoclonal Antibody - Images





Western blot analysis of HMGB2 expression in (1) HeLa cell lysate; (2) PC-12 cell lysate.