

**Anti-Cyclophilin F Rabbit Monoclonal Antibody**  
Catalog # ABO14457**Specification****Anti-Cyclophilin F Rabbit Monoclonal Antibody - Product Information**

Application	WB, IHC, IP
Primary Accession	<a href="#">P30405</a>
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

**Description**

Anti-Cyclophilin F Rabbit Monoclonal Antibody . Tested in WB, IHC, IP applications. This antibody reacts with Human, Mouse, Rat.

**Anti-Cyclophilin F Rabbit Monoclonal Antibody - Additional Information**

**Gene ID** 10105

**Other Names**

Peptidyl-prolyl cis-trans isomerase F, mitochondrial, PPIase F, 5.2.1.8, Cyclophilin D, CyP-D, CypD, Cyclophilin F, Mitochondrial cyclophilin, CyP-M, Rotamase F, PPIF, CYP3

**Calculated MW**

18 kDa KDa

**Application Details**

WB 1:500-1:2000<br>IHC 1:50-1:200<br>IP 1:50

**Contents**

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

**Immunogen**

A synthesized peptide derived from human Cyclophilin F

**Purification**

Affinity-chromatography

**Storage**

**Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.**

**Anti-Cyclophilin F Rabbit Monoclonal Antibody - Protein Information**

**Name** PPIF

## Synonyms CYP3

### Function

PPIase that catalyzes the cis-trans isomerization of proline imidic peptide bonds in oligopeptides and may therefore assist protein folding (PubMed: [20676357](http://www.uniprot.org/citations/20676357)). Involved in regulation of the mitochondrial permeability transition pore (mPTP) (PubMed: [26387735](http://www.uniprot.org/citations/26387735)). It is proposed that its association with the mPTP is masking a binding site for inhibiting inorganic phosphate (Pi) and promotes the open probability of the mPTP leading to apoptosis or necrosis; the requirement of the PPIase activity for this function is debated (PubMed: [26387735](http://www.uniprot.org/citations/26387735)). In cooperation with mitochondrial p53/TP53 is involved in activating oxidative stress-induced necrosis (PubMed: [22726440](http://www.uniprot.org/citations/22726440)). Involved in modulation of mitochondrial membrane F(1)F(0) ATP synthase activity and regulation of mitochondrial matrix adenine nucleotide levels (By similarity). Has anti-apoptotic activity independently of mPTP and in cooperation with BCL2 inhibits cytochrome c-dependent apoptosis (PubMed: [19228691](http://www.uniprot.org/citations/19228691)).

### Cellular Location

Mitochondrion matrix

## Anti-Cyclophilin F Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## Anti-Cyclophilin F Rabbit Monoclonal Antibody - Images




Figure 1. Western blot analysis of Cyclophilin F using anti-Cyclophilin F antibody (M02803). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human Hela whole cell lysates,

Lane 2: human HEL whole cell lysates,

Lane 3: human PC-3 whole cell lysates,

Lane 4: human HepG2 whole cell lysates,

Lane 5: rat PC-12 whole cell lysates,

Lane 6: rat C6 whole cell lysates,

Lane 7: mouse RAW264.7 whole cell lysates,

Lane 8: mouse Neuro-2a whole cell lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Cyclophilin F antigen affinity purified monoclonal antibody (Catalog # M02803) at 1:500 overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for Cyclophilin F at approximately 18 kDa. The expected band size for Cyclophilin F is at 22 kDa.