

## Anti-RAP80 Monoclonal Antibody Catalog # ABO14525

### Specification

---

#### Anti-RAP80 Monoclonal Antibody - Product Information

|                   |                        |
|-------------------|------------------------|
| Application       | WB, IHC, IF, ICC       |
| Primary Accession | <a href="#">Q96RL1</a> |
| Host              | Rabbit                 |
| Isotype           | Rabbit IgG             |
| Reactivity        | Human                  |
| Clonality         | Monoclonal             |
| Format            | Liquid                 |

#### Description

Anti-RAP80 Monoclonal Antibody . Tested in WB, IHC, ICC/IF applications. This antibody reacts with Human.

#### Anti-RAP80 Monoclonal Antibody - Additional Information

Gene ID 51720

#### Other Names

BRCA1-A complex subunit RAP80, Receptor-associated protein 80, Retinoid X receptor-interacting protein 110, Ubiquitin interaction motif-containing protein 1, UIMC1, RAP80, RXRIP110

#### Application Details

WB 1:1000-1:5000<br>IHC 1:50-1:200<br>ICC/IF 1:50-1:200

#### Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

#### Immunogen

A synthesized peptide derived from human RAP80 Ubiquitin-binding protein that specifically recognizes and binds 'Lys-63'-linked ubiquitin. Plays a central role in the BRCA1-A complex by specifically binding 'Lys-63'-linked ubiquitinated histones H2A and H2AX at DNA lesions sites, leading to target the BRCA1-BARD1 heterodimer to sites of DNA damage at double-strand breaks (DSBs).

#### Purification

Affinity-chromatography

Storage

**Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.**

#### Anti-RAP80 Monoclonal Antibody - Protein Information

**Name** UIMC1

**Synonyms** RAP80, RXRIP110

**Function**

Ubiquitin-binding protein (PubMed:<a href="http://www.uniprot.org/citations/24627472" target="\_blank">24627472</a>). Specifically recognizes and binds 'Lys-63'-linked ubiquitin (PubMed:<a href="http://www.uniprot.org/citations/19328070" target="\_blank">19328070</a>, Ref.38). Plays a central role in the BRCA1-A complex by specifically binding 'Lys-63'-linked ubiquitinated histones H2A and H2AX at DNA lesions sites, leading to target the BRCA1-BARD1 heterodimer to sites of DNA damage at double-strand breaks (DSBs). The BRCA1-A complex also possesses deubiquitinase activity that specifically removes 'Lys-63'- linked ubiquitin on histones H2A and H2AX. Also weakly binds monoubiquitin but with much less affinity than 'Lys-63'-linked ubiquitin. May interact with monoubiquitinated histones H2A and H2B; the relevance of such results is however unclear in vivo. Does not bind Lys-48'-linked ubiquitin. May indirectly act as a transcriptional repressor by inhibiting the interaction of NR6A1 with the corepressor NCOR1.

**Cellular Location**

Nucleus. Note=Localizes at sites of DNA damage at double-strand breaks (DSBs)

**Tissue Location**


Expressed in testis, ovary, thymus and heart. Expressed in germ cells of the testis.

**Anti-RAP80 Monoclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Anti-RAP80 Monoclonal Antibody - Images**



Western blot analysis of RAP80 expression in 293T cell lysate.