

Anti-PAK2 Monoclonal Antibody
Catalog # ABO14551

Specification

Anti-PAK2 Monoclonal Antibody - Product Information

Application	WB, IHC, IF, ICC, FC
Primary Accession	Q13177
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

Description

Anti-PAK2 Monoclonal Antibody . Tested in WB, IHC, ICC/IF, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

Anti-PAK2 Monoclonal Antibody - Additional Information

Gene ID 5062

Other Names

Serine/threonine-protein kinase PAK 2, 2.7.11.1, Gamma-PAK, PAK65, S6/H4 kinase, p21-activated kinase 2, PAK-2, p58, PAK-2p27, p27, PAK-2p34, p34, C-t-PAK2, PAK2

Application Details

WB 1:500-1:2000
IHC 1:100-1:500
ICC/IF 1:50-1:200
FC 1:80

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human PAK2 The activated kinase acts on a variety of targets. Phosphorylates ribosomal protein S6, histone H4 and myelin basic protein. Full length PAK 2 stimulates cell survival and cell growth. The process is, at least in part, mediated by phosphorylation and inhibition of pro-apoptotic BAD.

Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Anti-PAK2 Monoclonal Antibody - Protein Information

Name PAK2

Function

Serine/threonine protein kinase that plays a role in a variety of different signaling pathways including cytoskeleton regulation, cell motility, cell cycle progression, apoptosis or proliferation (PubMed: [12853446](http://www.uniprot.org/citations/12853446)), PubMed: [16617111](http://www.uniprot.org/citations/16617111)), PubMed: [19273597](http://www.uniprot.org/citations/19273597)), PubMed: [19923322](http://www.uniprot.org/citations/19923322)), PubMed: [33693784](http://www.uniprot.org/citations/33693784)), PubMed: [7744004](http://www.uniprot.org/citations/7744004)), PubMed: [9171063](http://www.uniprot.org/citations/9171063)). Acts as a downstream effector of the small GTPases CDC42 and RAC1 (PubMed: [7744004](http://www.uniprot.org/citations/7744004)). Activation by the binding of active CDC42 and RAC1 results in a conformational change and a subsequent autophosphorylation on several serine and/or threonine residues (PubMed: [7744004](http://www.uniprot.org/citations/7744004)). Full-length PAK2 stimulates cell survival and cell growth (PubMed: [7744004](http://www.uniprot.org/citations/7744004)). Phosphorylates MAPK4 and MAPK6 and activates the downstream target MAPKAPK5, a regulator of F-actin polymerization and cell migration (PubMed: [21317288](http://www.uniprot.org/citations/21317288)). Phosphorylates JUN and plays an important role in EGF-induced cell proliferation (PubMed: [21177766](http://www.uniprot.org/citations/21177766)). Phosphorylates many other substrates including histone H4 to promote assembly of H3.3 and H4 into nucleosomes, BAD, ribosomal protein S6, or MBP (PubMed: [21724829](http://www.uniprot.org/citations/21724829)). Phosphorylates CASP7, thereby preventing its activity (PubMed: [21555521](http://www.uniprot.org/citations/21555521)), PubMed: [27889207](http://www.uniprot.org/citations/27889207)). Additionally, associates with ARHGEF7 and GIT1 to perform kinase-independent functions such as spindle orientation control during mitosis (PubMed: [19273597](http://www.uniprot.org/citations/19273597)), PubMed: [19923322](http://www.uniprot.org/citations/19923322)). On the other hand, apoptotic stimuli such as DNA damage lead to caspase-mediated cleavage of PAK2, generating PAK-2p34, an active p34 fragment that translocates to the nucleus and promotes cellular apoptosis involving the JNK signaling pathway (PubMed: [12853446](http://www.uniprot.org/citations/12853446)), PubMed: [16617111](http://www.uniprot.org/citations/16617111)), PubMed: [9171063](http://www.uniprot.org/citations/9171063)). Caspase-activated PAK2 phosphorylates MKNK1 and reduces cellular translation (PubMed: [15234964](http://www.uniprot.org/citations/15234964)).

Cellular Location

[Serine/threonine-protein kinase PAK 2]: Cytoplasm Nucleus Note=MYO18A mediates the cellular distribution of the PAK2-ARHGEF7-GIT1 complex to the inner surface of the cell membrane

Tissue Location

Ubiquitously expressed. Higher levels seen in skeletal muscle, ovary, thymus and spleen


Anti-PAK2 Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)

- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-PAK2 Monoclonal Antibody - Images



Western blot analysis of PAK2 expression in HeLa cell lysate.