

Anti-TRPM7 Monoclonal Antibody Catalog # ABO14560

Specification

Anti-TRPM7 Monoclonal Antibody - Product Information

Application	WB, IF, ICC, FC
Primary Accession	Q96QT4
Host	Rabbit
Isotype	0.5-1mg/ml, actual concentration vary by lot. Use suggested dilution ratio to decide dilution procedure.
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

Description

Anti-TRPM7 Monoclonal Antibody . Tested in WB, ICC/IF, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

Anti-TRPM7 Monoclonal Antibody - Additional Information

Gene ID 54822

Other Names

Transient receptor potential cation channel subfamily M member 7, 2.7.11.1, Channel-kinase 1, Long transient receptor potential channel 7, LTrpC-7, LTrpC7, TRPM7, CHAK1, LTRPC7

Application Details

WB 1:500-1:2000
ICC/IF 1:50-1:200
FC 1:100

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human TRPM7

Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Anti-TRPM7 Monoclonal Antibody - Protein Information

Name TRPM7

Synonyms CHAK1, LTRPC7 {ECO:0000303|PubMed:113855}

Function

Bifunctional protein that combines an ion channel with an intrinsic kinase domain, enabling it to modulate cellular functions either by conducting ions through the pore or by phosphorylating downstream proteins via its kinase domain. The channel is highly permeable to divalent cations, specifically calcium (Ca²⁺), magnesium (Mg²⁺) and zinc (Zn²⁺) and mediates their influx (PubMed:11385574, PubMed:12887921, PubMed:15485879, PubMed:24316671, PubMed:35561741, PubMed:36027648). Controls a wide range of biological processes such as Ca²⁺(+), Mg²⁺(+) and Zn²⁺(+) homeostasis, vesicular Zn²⁺ release channel and intracellular Ca²⁺ signaling, embryonic development, immune responses, cell motility, proliferation and differentiation (By similarity). The C-terminal alpha-kinase domain autophosphorylates cytoplasmic residues of TRPM7 (PubMed:18365021). In vivo, TRPM7 phosphorylates SMAD2, suggesting that TRPM7 kinase may play a role in activating SMAD signaling pathways. In vitro, TRPM7 kinase phosphorylates ANXA1 (annexin A1), myosin II isoforms and a variety of proteins with diverse cellular functions (PubMed:15485879, PubMed:18394644).

Cellular Location

Cell membrane; Multi-pass membrane protein {ECO:0000250|UniProtKB:Q923J1}. Cytoplasmic vesicle membrane {ECO:0000250|UniProtKB:Q923J1}; Multi-pass membrane protein {ECO:0000250|UniProtKB:Q923J1}. Note=Localized largely in intracellular Zn²⁺-storage vesicles. {ECO:0000250|UniProtKB:Q923J1}


Anti-TRPM7 Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-TRPM7 Monoclonal Antibody - Images





Western blot analysis of TRPM7 expression in HeLa cell lysate.