

Anti-KMT5A / SETD8 / Pr-SET7 Monoclonal Antibody
Catalog # ABO14566**Specification****Anti-KMT5A / SETD8 / Pr-SET7 Monoclonal Antibody - Product Information**

Application	WB, FC
Primary Accession	O9NOR1
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

Description

Anti-KMT5A / SETD8 / Pr-SET7 Monoclonal Antibody . Tested in WB, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

Anti-KMT5A / SETD8 / Pr-SET7 Monoclonal Antibody - Additional Information

Gene ID 387893

Other Names

N-lysine methyltransferase KMT5A, 2.1.1.-, H4-K20-HMTase KMT5A, Histone-lysine N-methyltransferase KMT5A, 2.1.1.361, Lysine N-methyltransferase 5A, Lysine-specific methylase 5A {ECO:0000312|HGNC:HGNC:29489}, PR/SET domain-containing protein 07, PR-Set7, PR/SET07, SET domain-containing protein 8, KMT5A (http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=29489)

Application Details

WB 1:1000-1:5000
FC 1:100

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human KMT5A / SETD8 / Pr-SET7 Protein-lysine N-methyltransferase that monomethylates both histones and non-histone proteins. Specifically monomethylates 'Lys-20' of histone H4 (H4K20me1). H4K20me1 is enriched during mitosis and represents a specific tag for epigenetic transcriptional repression. Mainly functions in euchromatin regions, thereby playing a central role in the silencing of euchromatic genes.

Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Anti-KMT5A / SETD8 / Pr-SET7 Monoclonal Antibody - Protein Information

Name KMT5A ([HGNC:29489](#))

Function

Protein-lysine N-methyltransferase that monomethylates both histones and non-histone proteins (PubMed: [12086618](#), PubMed: [12121615](#), PubMed: [15964846](#), PubMed: [17707234](#), PubMed: [27338793](#)). Specifically monomethylates 'Lys-20' of histone H4 (H4K20me1) (PubMed: [12086618](#), PubMed: [12121615](#), PubMed: [15200950](#), PubMed: [15933069](#), PubMed: [15933070](#), PubMed: [15964846](#), PubMed: [16517599](#), PubMed: [27338793](#)). H4K20me1 is enriched during mitosis and represents a specific tag for epigenetic transcriptional repression (PubMed: [12086618](#), PubMed: [12121615](#), PubMed: [15200950](#), PubMed: [15933069](#), PubMed: [15933070](#), PubMed: [15964846](#), PubMed: [16517599](#)). Mainly functions in euchromatin regions, thereby playing a central role in the silencing of euchromatic genes (PubMed: [12086618](#), PubMed: [12121615](#), PubMed: [15200950](#), PubMed: [15933069](#), PubMed: [15933070](#), PubMed: [15964846](#), PubMed: [16517599](#)). Required for cell proliferation, probably by contributing to the maintenance of proper higher-order structure of DNA during mitosis (PubMed: [12086618](#), PubMed: [12121615](#), PubMed: [15200950](#), PubMed: [15933069](#), PubMed: [15933070](#), PubMed: [15964846](#), PubMed: [16517599](#)). Involved in chromosome condensation and proper cytokinesis (PubMed: [12086618](#), PubMed: [12121615](#), PubMed: [15200950](#), PubMed: [15933069](#), PubMed: [15933070](#), PubMed: [15964846](#), PubMed: [16517599](#)). Nucleosomes are preferred as substrate compared to free histones (PubMed: [12086618](#), PubMed: [12121615](#), PubMed: [15200950](#), PubMed: [15933069](#), PubMed: [15933070](#), PubMed: [15964846](#), PubMed: [16517599](#)).

href="http://www.uniprot.org/citations/12121615" target="_blank">12121615, PubMed:15200950, PubMed:15933069, PubMed:15933070, PubMed:15964846, PubMed:16517599). Mediates monomethylation of p53/TP53 at 'Lys-382', leading to repress p53/TP53-target genes (PubMed:17707234). Plays a negative role in TGF- beta response regulation and a positive role in cell migration (PubMed:23478445).

Cellular Location


Nucleus. Chromosome. Note=Specifically localizes to mitotic chromosomes (PubMed:12208845). Colocalized with SIRT2 at mitotic foci (PubMed:23468428). Associates with chromosomes during mitosis; association is increased in a H₂O₂-induced oxidative stress- dependent manner (PubMed:23468428). Associates with silent chromatin on euchromatic arms (PubMed:12086618). Not associated with constitutive heterochromatin (PubMed:12086618).

Anti-KMT5A / SETD8 / Pr-SET7 Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-KMT5A / SETD8 / Pr-SET7 Monoclonal Antibody - Images



Western blot analysis of KMT5A / SETD8 / Pr-SET7 expression in (1) 293T cell lysate; (2) NIH/3T3 cell lysate.