

Anti-CD7 Monoclonal Antibody
Catalog # ABO14583**Specification****Anti-CD7 Monoclonal Antibody - Product Information**

Application	WB, IHC, IF, ICC, IP
Primary Accession	P09564
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid

Description

Anti-CD7 Monoclonal Antibody . Tested in WB, IHC, ICC/IF, IP applications. This antibody reacts with Human.

Anti-CD7 Monoclonal Antibody - Additional Information

Gene ID 924

Other Names

T-cell antigen CD7, GP40, T-cell leukemia antigen, T-cell surface antigen Leu-9, TP41, CD7, CD7

Application Details

WB 1:500-1:2000
IHC 1:50-1:200
ICC/IF 1:50-1:200
IP 1:50

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human CD7 Not yet known.

Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Anti-CD7 Monoclonal Antibody - Protein Information

Name CD7

Function

Transmembrane glycoprotein expressed by T-cells and natural killer (NK) cells and their precursors (PubMed:7506726).

Plays a costimulatory role in T-cell activation upon binding to its ligand K12/SECTM1 (PubMed:10652336). In turn, mediates the production of cytokines such as IL-2 (PubMed:1709867). On resting NK-cells, CD7 activation results in a significant induction of interferon-gamma levels (PubMed:7506726).

Cellular Location

Membrane; Single-pass type I membrane protein.

Tissue Location


Expressed on T-cells and natural killer (NK) cells and their precursors.

Anti-CD7 Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-CD7 Monoclonal Antibody - Images



Western blot analysis of CD7 expression in human PBMC lysate.