

## Anti-VPS35 Monoclonal Antibody Catalog # ABO14625

### Specification

---

#### Anti-VPS35 Monoclonal Antibody - Product Information

Application	WB, IHC, IF, ICC, FC
Primary Accession	<a href="#">O96QK1</a>
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

#### Description

Anti-VPS35 Monoclonal Antibody . Tested in WB, IHC, ICC/IF, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

#### Anti-VPS35 Monoclonal Antibody - Additional Information

Gene ID 55737

#### Other Names

Vacuolar protein sorting-associated protein 35, hVPS35, Maternal-embryonic 3, Vesicle protein sorting 35, VPS35 {ECO:0000303|PubMed:28397838, ECO:0000312|HGNC:HGNC:13487}

#### Application Details

WB 1:1000-1:5000<br>IHC 1:50-1:200<br>ICC/IF 1:50-1:200<br>FC 1:50

#### Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

#### Immunogen

A synthesized peptide derived from human VPS35

#### Purification

Affinity-chromatography

Storage

**Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.**

#### Anti-VPS35 Monoclonal Antibody - Protein Information

**Name** VPS35 {ECO:0000303|PubMed:28397838, ECO:0000312|HGNC:HGNC:13487}

#### Function

Acts as a component of the retromer cargo-selective complex (CSC). The CSC is believed to be the

core functional component of retromer or respective retromer complex variants acting to prevent missorting of selected transmembrane cargo proteins into the lysosomal degradation pathway. The recruitment of the CSC to the endosomal membrane involves RAB7A and SNX3. The CSC seems to associate with the cytoplasmic domain of cargo proteins predominantly via VPS35; however, these interactions seem to be of low affinity and retromer SNX proteins may also contribute to cargo selectivity thus questioning the classical function of the CSC. The SNX-BAR retromer mediates retrograde transport of cargo proteins from endosomes to the trans-Golgi network (TGN) and is involved in endosome-to-plasma membrane transport for cargo protein recycling. The SNX3-retromer mediates the retrograde endosome-to-TGN transport of WLS distinct from the SNX-BAR retromer pathway (PubMed:<a href="http://www.uniprot.org/citations/30213940" target="\_blank">30213940</a>). The SNX27-retromer is believed to be involved in endosome-to-plasma membrane trafficking and recycling of a broad spectrum of cargo proteins. The CSC seems to act as recruitment hub for other proteins, such as the WASH complex and TBC1D5 (Probable). Required for retrograde transport of lysosomal enzyme receptor IGF2R and SLC11A2. Required to regulate transcytosis of the polymeric immunoglobulin receptor (pIgR-pIgA) (PubMed:<a href="http://www.uniprot.org/citations/15078903" target="\_blank">15078903</a>, PubMed:<a href="http://www.uniprot.org/citations/15247922" target="\_blank">15247922</a>, PubMed:<a href="http://www.uniprot.org/citations/20164305" target="\_blank">20164305</a>). Required for endosomal localization of WASHC2C (PubMed:<a href="http://www.uniprot.org/citations/22070227" target="\_blank">22070227</a>, PubMed:<a href="http://www.uniprot.org/citations/28892079" target="\_blank">28892079</a>). Mediates the association of the CSC with the WASH complex via WASHC2 (PubMed:<a href="http://www.uniprot.org/citations/22070227" target="\_blank">22070227</a>, PubMed:<a href="http://www.uniprot.org/citations/24819384" target="\_blank">24819384</a>, PubMed:<a href="http://www.uniprot.org/citations/24980502" target="\_blank">24980502</a>). Required for the endosomal localization of TBC1D5 (PubMed:<a href="http://www.uniprot.org/citations/20923837" target="\_blank">20923837</a>).

#### Cellular Location

Cytoplasm. Membrane; Peripheral membrane protein. Endosome Early endosome. Late endosome  
Note=Localizes to tubular profiles adjacent to endosomes

#### Tissue Location

Ubiquitous. Highly expressed in heart, brain, placenta, skeletal muscle, spleen, thymus, testis, ovary, small intestine, kidney and colon


### Anti-VPS35 Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Anti-VPS35 Monoclonal Antibody - Images





Western blot analysis of VPS35 expression in (1) HeLa cell lysate; (2) Mouse kidney lysate.