

**Anti-CD10 MME Monoclonal Antibody**  
Catalog # ABO14758

**Specification**

---

**Anti-CD10 MME Monoclonal Antibody - Product Information**

Application	WB, IHC, IP
Primary Accession	<a href="#">P08473</a>
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human
Clonality	Monoclonal
Format	Liquid

**Description**

Anti-CD10 MME Monoclonal Antibody . Tested in WB, IHC, IP applications. This antibody reacts with Human, Rat.

**Anti-CD10 MME Monoclonal Antibody - Additional Information**

**Gene ID** 4311

**Other Names**

Neprilysin, 3.4.24.11, Atriopeptidase, Common acute lymphocytic leukemia antigen, CALLA, Enkephalinase, Neutral endopeptidase 24.11, NEP, Neutral endopeptidase, Skin fibroblast elastase, SFE, CD10, MME {ECO:0000303|PubMed:27588448, ECO:0000312|HGNC:HGNC:7154}

**Application Details**

WB 1:500-1:2000<br>IHC 1:50-1:200<br>IP 1:50

**Contents**

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

**Immunogen**

A synthesized peptide derived from human CD10 CD10 is a transmembrane type II molecule and functions as a metallo-peptidase requiring zinc. Specifically, CD10 cleaves 1-3 amino-terminal amino acids from peptides with a preference for neutral amino acids (valine, iso-leucine, phenylalanine, leucine or alanine). Involved in the degradation of atrial natriuretic factor (ANF).

**Purification**

Affinity-chromatography

**Storage**

**Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.**

**Anti-CD10 MME Monoclonal Antibody - Protein Information**

**Name** MME {ECO:0000303|PubMed:27588448, ECO:0000312|HGNC:HGNC:7154}

### Function

Thermolysin-like specificity, but is almost confined on acting on polypeptides of up to 30 amino acids (PubMed:<a href="http://www.uniprot.org/citations/15283675" target="\_blank">15283675</a>, PubMed:<a href="http://www.uniprot.org/citations/6208535" target="\_blank">6208535</a>, PubMed:<a href="http://www.uniprot.org/citations/6349683" target="\_blank">6349683</a>, PubMed:<a href="http://www.uniprot.org/citations/8168535" target="\_blank">8168535</a>). Biologically important in the destruction of opioid peptides such as Met- and Leu-enkephalins by cleavage of a Gly-Phe bond (PubMed:<a href="http://www.uniprot.org/citations/17101991" target="\_blank">17101991</a>, PubMed:<a href="http://www.uniprot.org/citations/6349683" target="\_blank">6349683</a>). Catalyzes cleavage of bradykinin, substance P and neurotensin peptides (PubMed:<a href="http://www.uniprot.org/citations/6208535" target="\_blank">6208535</a>). Able to cleave angiotensin-1, angiotensin-2 and angiotensin 1-9 (PubMed:<a href="http://www.uniprot.org/citations/15283675" target="\_blank">15283675</a>, PubMed:<a href="http://www.uniprot.org/citations/6349683" target="\_blank">6349683</a>). Involved in the degradation of atrial natriuretic factor (ANF) and brain natriuretic factor (BNP(1-32)) (PubMed:<a href="http://www.uniprot.org/citations/16254193" target="\_blank">16254193</a>, PubMed:<a href="http://www.uniprot.org/citations/2531377" target="\_blank">2531377</a>, PubMed:<a href="http://www.uniprot.org/citations/2972276" target="\_blank">2972276</a>). Displays UV-inducible elastase activity toward skin preelastic and elastic fibers (PubMed:<a href="http://www.uniprot.org/citations/20876573" target="\_blank">20876573</a>).

### Cellular Location

Cell membrane; Single-pass type II membrane protein


### Anti-CD10 MME Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Anti-CD10 MME Monoclonal Antibody - Images





Western blot analysis of CD10 expression in Ramos cell lysate.