

Anti-ARSA Antibody Picoband™ (monoclonal, 4C10)
Catalog # ABO14785

Specification

Anti-ARSA Antibody Picoband™ (monoclonal, 4C10) - Product Information

Application	WB, IHC, FC
Primary Accession	P15289
Host	Mouse
Isotype	Mouse IgG2a
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Lyophilized

Description

Anti-ARSA Antibody Picoband™ (monoclonal, 4C10) . Tested in Flow Cytometry, IHC, WB applications. This antibody reacts with Human, Mouse, Rat.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-ARSA Antibody Picoband™ (monoclonal, 4C10) - Additional Information

Gene ID 410

Other Names

Arylsulfatase A, ASA, 3.1.6.8, Cerebroside-sulfatase, Arylsulfatase A component B, Arylsulfatase A component C, ARSA

Calculated MW

54 kDa KDa

Application Details

Western blot, 0.1-0.5 µg/ml, Human, Mouse, Rat
 Immunohistochemistry (Paraffin-embedded Section), 0.5-1 µg/ml, Human, By Heat
 Flow Cytometry, 1-3 µg/1x10⁶ cells, Human, Mouse

Subcellular Localization

Lysosome.

Contents

Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg Na₃N.

Immunogen

A synthetic peptide corresponding to a sequence at the C-terminus of human ARSA, different from the related mouse sequence by six amino acids.

Cross Reactivity

No cross-reactivity with other proteins.

Storage

Store at -20°C for one year from date of

receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.

Anti-ARSA Antibody Picoband™ (monoclonal, 4C10) - Protein Information

Name ARSA

Function

Hydrolyzes cerebroside sulfate.

Cellular Location

Endoplasmic reticulum. Lysosome

Anti-ARSA Antibody Picoband™ (monoclonal, 4C10) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-ARSA Antibody Picoband™ (monoclonal, 4C10) - Images




Figure 2. Western blot analysis of ARSA using anti-ARSA antibody (M02583). Electrophoresis was performed on a 10% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions.

Lane 1: rat testis tissue lysate,
Lane 2: rat liver tissue lysate,
Lane 3: rat brain tissue lysate,
Lane 4: rat lung tissue lysate,
Lane 5: mouse testis tissue lysate,
Lane 6: mouse liver tissue lysate,
Lane 7: mouse brain tissue lysate,

Lane 8: mouse lung tissue lysate,
 Lane 9: mouse HEPA1-6 whole cell lysate.

After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with mouse anti-ARSA antigen affinity purified monoclonal antibody (Catalog # M02583) at 0.5 µg/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-mouse IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1001) with Tanon 5200 system.




Figure 3. Western blot analysis of ARSA using anti-ARSA antibody (M02583).

Electrophoresis was performed on a 10% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50µg of sample under reducing conditions.

Lane 1: human A375 whole cell lysate,
 Lane 2: human A549 whole cell lysate,
 Lane 3: human SMMC-7721 whole cell lysate.

After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with mouse anti-ARSA antigen affinity purified monoclonal antibody (Catalog # M02583) at 0.5 µg/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-mouse IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1001) with Tanon 5200 system.




Figure 1. Flow Cytometry analysis of ANA-1 cells using anti-ARSA antibody (M02583).

Overlay histogram showing ANA-1 cells stained with M02583 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with mouse anti-ARSA Antibody (M02583, 1 $\mu\text{g}/1 \times 10^6$ cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-mouse IgG (BA1126, 5-10 $\mu\text{g}/1 \times 10^6$ cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was mouse IgG (1 $\mu\text{g}/1 \times 10^6$) used under the same conditions. Unlabelled sample (Red line) was also used as a control.




Figure 4. Flow Cytometry analysis of Raji cells using anti-ARSA antibody (M02583).

Overlay histogram showing Raji cells stained with M02583 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-ARSA Antibody (M02583, 1 $\mu\text{g}/1 \times 10^6$ cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-rabbit IgG (BA1127, 5-10 $\mu\text{g}/1 \times 10^6$ cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG (1 $\mu\text{g}/1 \times 10^6$) used under the same conditions. Unlabelled sample (Red line) was also used as a control.




Figure 5. IHC analysis of ARSA using anti-ARSA antibody (M02583).

ARSA was detected in paraffin-embedded section of human lung cancer tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1 $\mu\text{g}/\text{ml}$ rabbit anti-ARSA Antibody (M02583) overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC)(Catalog # SA1022) with DAB as the chromogen.

Anti-ARSA Antibody Picoband™ (monoclonal, 4C10) - Background

Arylsulfatase A (ARSA) is an enzyme that breaks down sulfatides, namely cerebroside 3-sulfate into cerebroside and sulfate. In humans, arylsulfatase A is encoded by the ARSA gene. ARSA is mapped to 22q13.33. The protein encoded by this gene hydrolyzes cerebroside sulfate to

cerebroside and sulfate. Defects in this gene lead to metachromatic leucodystrophy (MLD), a progressive demyelination disease which results in a variety of neurological symptoms and ultimately death. Alternatively spliced transcript variants have been described for this gene.