

Anti-CASR Antibody Picoband™ (monoclonal, 11E9)
Catalog # ABO14786

Specification

Anti-CASR Antibody Picoband™ (monoclonal, 11E9) - Product Information

Application	WB, IHC, ICC, FC
Primary Accession	P41180
Host	Mouse
Isotype	Mouse IgG2b
Reactivity	Human
Clonality	Monoclonal
Format	Lyophilized

Description

Anti-CASR Antibody Picoband™ (monoclonal, 11E9) . Tested in Flow Cytometry, IHC, ICC, WB applications. This antibody reacts with Human.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-CASR Antibody Picoband™ (monoclonal, 11E9) - Additional Information

Gene ID 846

Other Names

Extracellular calcium-sensing receptor, CaR, CaSR, hCasR, Parathyroid cell calcium-sensing receptor 1, PCaR1, CASR ([HGNC:1514](http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=1514))

Calculated MW

130 kDa KDa

Application Details

Western blot, 0.1-0.5 µg/ml
Immunohistochemistry (Paraffin-embedded Section), 0.5-1 µg/ml
Immunohistochemistry (Frozen Section), 0.5-1 µg/ml
Immunocytochemistry, 0.5-1 µg/ml
Flow Cytometry, 1-3 µg/1x10⁶ cells

Subcellular Localization

Cell membrane

Tissue Specificity

Expressed in the temporal lobe, frontal lobe, parietal lobe, hippocampus, and cerebellum. Also found in kidney, lung, liver, heart, skeletal muscle, placenta.

Contents

Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg Na₃N.

Immunogen

E. coli-derived human CASR recombinant protein (Position: Q926-S1078). Human CASR shares

80.5% and 78.6% amino acid (aa) sequence identity with mouse and rat CASR, respectively.

Cross Reactivity

No cross-reactivity with other proteins.

Storage

Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.

Anti-CASR Antibody Picoband™ (monoclonal, 11E9) - Protein Information

Name CASR ([HGNC:1514](#))

Function

G-protein-coupled receptor that senses changes in the extracellular concentration of calcium ions and plays a key role in maintaining calcium homeostasis (PubMed:[17555508](http://www.uniprot.org/citations/17555508), PubMed:[19789209](http://www.uniprot.org/citations/19789209), PubMed:[21566075](http://www.uniprot.org/citations/21566075), PubMed:[22114145](http://www.uniprot.org/citations/22114145), PubMed:[22789683](http://www.uniprot.org/citations/22789683), PubMed:[23966241](http://www.uniprot.org/citations/23966241), PubMed:[25104082](http://www.uniprot.org/citations/25104082), PubMed:[25292184](http://www.uniprot.org/citations/25292184), PubMed:[25766501](http://www.uniprot.org/citations/25766501), PubMed:[26386835](http://www.uniprot.org/citations/26386835), PubMed:[7759551](http://www.uniprot.org/citations/7759551), PubMed:[8636323](http://www.uniprot.org/citations/8636323), PubMed:[8702647](http://www.uniprot.org/citations/8702647), PubMed:[8878438](http://www.uniprot.org/citations/8878438)). Senses fluctuations in the circulating calcium concentration and modulates the production of parathyroid hormone (PTH) in parathyroid glands (By similarity). The activity of this receptor is mediated by a G-protein that activates a phosphatidylinositol-calcium second messenger system (PubMed:[7759551](http://www.uniprot.org/citations/7759551)). The G-protein-coupled receptor activity is activated by a co-agonist mechanism: aromatic amino acids, such as Trp or Phe, act concertedly with divalent cations, such as calcium or magnesium, to achieve full receptor activation (PubMed:[27386547](http://www.uniprot.org/citations/27386547), PubMed:[27434672](http://www.uniprot.org/citations/27434672)).

Cellular Location

Cell membrane; Multi-pass membrane protein

Tissue Location

Expressed in the temporal lobe, frontal lobe, parietal lobe, hippocampus, and cerebellum. Also found in kidney, lung, liver, heart, skeletal muscle, placenta.

Anti-CASR Antibody Picoband™ (monoclonal, 11E9) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)

- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-CASR Antibody Picoband™ (monoclonal, 11E9) - Images




Figure 2. Western blot analysis of CASR using anti-CASR antibody (M00574). Electrophoresis was performed on a 8% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions.

- Lane 1: human Hela whole cell lysate,
- Lane 2: human A549 whole cell lysate,
- Lane 3: human 22RV1 whole cell lysate,
- Lane 4: human HepG2 whole cell lysate,
- Lane 5: human Caco-2 whole cell lysate.

After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with mouse anti-CASR antigen affinity purified monoclonal antibody (Catalog # M00574) at 0.5 µg/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-mouse IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1001) with Tanon 5200 system.




Figure 1. Flow Cytometry analysis of NEURO-2A cells using anti-CASR antibody (M00574).

Overlay histogram showing NEURO-2A cells stained with M00574 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with mouse anti-CASR Antibody (M00574, 1 $\mu\text{g}/1 \times 10^6$ cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-mouse IgG (BA1126, 5-10 $\mu\text{g}/1 \times 10^6$ cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was mouse IgG (1 $\mu\text{g}/1 \times 10^6$) used under the same conditions. Unlabelled sample (Red line) was also used as a control.

Anti-CASR Antibody Picoband™ (monoclonal, 11E9) - Background

The calcium-sensing receptor (CaSR) is a G protein-coupled receptor that is expressed in the parathyroid hormone (PTH)-producing chief cells of the parathyroid gland, and the cells lining the kidney tubule. It senses small changes in circulating calcium concentration and couples this information to intracellular signaling pathways that modify PTH secretion or renal cation handling, thus this protein plays an essential role in maintaining mineral ion homeostasis. Mutations in this gene cause familial hypocalciuric hypercalcemia, familial, isolated hypoparathyroidism, and neonatal severe primary hyperparathyroidism.